



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-Fluorescein [4-4-20 (enhanced)] Standard Size Ab00102-10.6

This is a Fab fragment with a his-tag.

This is a chimeric antibody created as part of a panel offering antibodies of the same specificity in different formats (species, isotype, subtype and modified versions) for use as isotype controls.

**Isotype and Format:** Human Fab fragment, His-Tagged, Kappa

**Clone Number:** 4-4-20 (enhanced)

**Alternative Name(s) of Target:** FI; fluorescyl; fluoresceine; FITC

**UniProt Accession Number of Target Protein:** n/a

**Published Application(s):** negative control, WB, FC

**Published Species Reactivity:** n/a

**Immunogen:** Fluorescein I-bovine serum albumin (FII-BSA).

**Specificity:** Fluorescein labelled proteins.

**Application Notes:**

**Antibody First Published in:** Jung S, Honegger A, Plückthun A. Selection for improved protein stability by phage display. J Mol Biol. 1999 Nov 19;294(1):163-80. [PMID:10556036](#)

**Note on publication:** Describes the use of phage display to improve 4-4-20 variable domain, including a 20-fold better binding constant.

### Product Form

**Size:** 200 µg Purified antibody.

**Purification:** Purified by Immobilized Metal Affinity Chromatography

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.