



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Fluorescein [4-4-20 (enhanced)] Standard Size Ab00102-7.4

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This is a chimeric antibody created as part of a panel offering antibodies of the same specificity in different formats (species, isotype, subtype and modified versions) for use as isotype controls.

Isotype and Format: Rat IgG2a, Fc Silent™, Kappa

Clone Number: 4-4-20 (enhanced)

Alternative Name(s) of Target: Fl; fluorescyl; fluoresceine; FITC

UniProt Accession Number of Target Protein: n/a

Published Application(s): negative control, WB, FC

Published Species Reactivity: n/a

Immunogen: Fluorescein I-bovine serum albumin (FII-BSA).

Specificity: Fluorescein labelled proteins.

Application Notes:

Antibody First Published in: Jung S, Honegger A, Plückthun A. Selection for improved protein stability by phage display. J Mol Biol. 1999 Nov 19;294(1):163-80. [PMID:10556036](#)

Note on publication: Describes the use of phage display to improve 4-4-20 variable domain, including a 20-fold better binding constant.

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.