



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Thrombopoietin [TN1] Bulk Size Ab00746-23.0-BT

This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG1 format, for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Rabbit IgG, Kappa

Clone Number: TN1

Alternative Name(s) of Target: C-mpl ligand; Megakaryocyte colony-stimulating factor; Megakaryocyte growth and development factor; TPO

UniProt Accession Number of Target Protein: P40225

Published Application(s): crystallization, neutralize, ELISA

Published Species Reactivity: Human

Immunogen: TN1 was prepared by immunizing BALB/c mice by intraperitoneal injection of recombinant human TPO. The Fab fragment was produced via incubation with papain which cleaves at the hinge region of the IgG to remove the Fc region. The product is then subsequently incubated under reducing conditions and the free Cys residue blocked with iodoacetamide.

Specificity: TN1 binds specifically to TPO at an epitope which consists of residues from the C-terminal end of helix C (R98) and the C-D crossover loop (residues 102-115) and residues from the C-terminal half of helix B (residues 57-61) (PMID: 14769915). TPO is a heavily glycosylated cytokine which acts as the ligand for the hematopoietic receptor c-Mpl. The primary role of TPO is as a regulator of megakaryocytopoiesis and platelet production. Stimulated c-Mpl transduces signals into target cells via the JAK/STAT, Ras and MAPK pathways.

Application Notes: TN1 can neutralize the biological activity of TPO and can be used to detect TPO in ELISA studies. The Fab fragment of TN1 can also be used to aid the crystallisation of TPO by increasing the area upon which stable crystal contacts can be made (PMID: 11976502).

Antibody First Published in: Tahara et al. A sensitive sandwich ELISA for measuring thrombopoietin in human serum: serum thrombopoietin levels in healthy volunteers and in patients with haemopoietic disorders. British Journal of Haematology 1996, 93, 783-788 [PMID:](#)

Note on publication: Describes the use of TN1 in the development of a reproducible, highly sensitive and specific human TPO ELISA procedure

Product Form

Size:

1 mg Purified antibody in bulk size.

Purification: Protein A affinity purified

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg /ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.