

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Anti-trans-sialidase [13G9] Standard Size Ab00762-2.0

Isotype and Format: Mouse IgG2a, Kappa

Clone Number: 13G9

Alternative Name(s) of Target: TS

UniProt Accession Number of Target Protein: Q26966 Published Application(s): crystallization, neutralize, IF Published Species Reactivity: Trypanosoma cruzi

Immunogen: 13G9 was prepared by immnuizing mice with recombinant TS (Δ 1443TS), which is fully active but includes a deletion of a non-neutralizing epitope (speeds up production of neutralizing antibodies). The antibody was selected for by screening using a TS-inhibition assay.

Specificity: 13G9 is a neutralizing antibody which binds specifically and with high affinity (KD ~0.72 nM) to a conformational epitope on the Trypanosoma cruzi TS protein – the antibody binds to TS in such a way as to prevent the movement of Y119, whose mobility has a key role in the trans-glycosidase mechanism. Specificity of binding was deduced by the absence of reactivity against TS from the closely related Trypanosoma rangeli and Trypanosoma brucei. A Fab fragment of the antibody was generated by digestion with papain - the Fab fragment possesses lower inhibitory activity against T. cruzi TS (Fab IC50 ~1.6nM; full-length mAb IC50 ~56pM). Trypanosoma cruzi is a protozoan parasite which causes Chagas' disease (or American trypanosomiasis) and its protein TS is a glycosylphosphatidylinositol-anchored non-integral membrane protein and a virulence factor which is involved in host cell invasion/survival, as well as being actively shed in the bloodstream. TS performs a glyosyl-transfer reaction to convert host glycoconjugates into sialic acids (cannot perform de novo synthesis of sialic acids, which are important for life cycle and survival).

Application Notes: 13G9 can be used for IF with whole parasites which allows membrane localization of TS to be observed. The antibody can inhibit TS-mediated sialylation of the parasite via the transfer of sialic acid from the enivronment to the surface of the parasite. Cell invasion (a process in which TS plays a key role) is also inhibited by 13G9 and was shown to greatly reduce the number of T. cruzi-infected cells. Passive transfer of 13G9 to T. cruzi-infected mice provided protection against the delterious effects that TS has on the immune system and platelets. The 13G9 Fab fragment was also used to crystallize TS for structure studies and this allowed the binding to TS to be seen.

Antibody First Published in: Buschiazzo et al. Trypanosoma cruzi trans-sialidase in complex with a neutralizing antibody: structure/function studies towards the rational design of inhibitors. PLoS Pathog. 2012 Jan;8(1):e1002474. PMID:

Note on publication: Describes the production of 13G9 and functional and structural studies on th mAb.

Product Form

Size: 200 µg Purified antibody.

Purification:

Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.