



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-Sag1 [4F11E12] Standard Size Ab00763-23.0

This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG1 format, for improved compatibility with existing reagents, assays and techniques.

**Isotype and Format:** Rabbit IgG, Kappa

**Clone Number:** 4F11E12

**Alternative Name(s) of Target:** P30

**UniProt Accession Number of Target Protein:** C7E5U4

**Published Application(s):** crystallization

**Published Species Reactivity:** Toxoplasma gondii

**Immunogen:** 4F11E12 was generated by intraperitoneal infection of mice with cysts from the 76 K T. gondii strain and was selected for its ability to inhibit efficiently the recognition of SAG1 by sera from infected patients. The 4F11E12 Fab fragment was created by digestion with papain

**Specificity:** 4F11E12 recognises a highly conserved conformational epitope on the surface of the D1 domain in both native and recombinant SAG1, located within the N-terminal domain; the binding site does not overlap with the proposed ligand-binding pocket or homodimerization site. The epitope is an immunodominant area and includes residues from 2 long loops located at one extremity. SAG1 is a GPI-anchored cell surface protein predicted to act as an adhesin during host-cell attachment through its binding to proteoglycans. SAG1 recognizes ligands such as glucosamide moieties, which are expressed in a broad range of host cell types. T. gondii is a protozoan parasite capable of causing chronic infections resulting in congenital disease and has the ability to evade natural host defences.

**Application Notes:** F411E12 can be co-crystallized with SAG1, which enables the mode of binding to be elucidated.

**Antibody First Published in:** Graille et al. Crystal structure of the complex between the monomeric form of Toxoplasma gondii surface antigen 1 (SAG1) and a monoclonal antibody that mimics the human immune response. J Mol Biol. 2005 Nov 25;354(2):447-58. Epub 2005 Sep 30. [PMID:16242717](#)

**Note on publication:** Describes the generation of F411E12 Fab and its co-crystallization with SAG1. The epitope binding site was determined and it was found that neither homodimerization or prevents access to the ligand binding site (which are necessary to neutralize parasite entry into host cells).

## Product Form

**Size:**

200 µg Purified antibody.

**Purification:** Protein A affinity purified

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.