



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-alpha-2 integrin inserted-domain [2E10] Standard Size Ab00777-1.1

This reformatted mouse antibody was made using the variable domain sequences of the original Mouse IgG (scFv) format, for improved compatibility with existing reagents, assays and techniques.

**Isotype and Format:** Mouse IgG1, Kappa

**Clone Number:** 2E10

**Alternative Name(s) of Target:** alpha2-beta1 integrin; CD49b; glycoprotein Ialla; GP Ialla; VLA2; VLA-2; collagen receptor; alpha-2 integrin I-domain; integrin alpha-2; integrin alpha 2

**UniProt Accession Number of Target Protein:** P17301

**Published Application(s):** ELISA, FC

**Published Species Reactivity:** Human

**Immunogen:** 2E10 was generated by immunizing BALB/c mice with a DNA vaccine containing the human alpha2 I-domain cDNA sequence within pVAC2. The alpha2 I-domain is expressed as a fusion protein with fragment C of tetanus toxoid (FrC-TT). The scFv was then created using a V-gene phage display library derived from RNA from mice splenocytes and isolated following selection on GST-I after 3 rounds of panning.

**Specificity:** scFv 2E10 can bind to the I (inserted) domain of recombinant alpha2 (GST- and His-), but cannot bind to the native epitope on platelets. This scFv also cross-reacts with and binds to the I-domain of alpha1 integrin. The alpha2 I-domain consists of ~200 amino acids and contains the ligand-binding site. The  $\alpha 2\beta 1$  integrin is expressed on platelets, fibroblasts, activated T lymphocytes, endothelial cells and epithelial cells where its function is to act as a collagen and laminin receptor.

**Application Notes:** 2E10 scFv can be used to detect alpha2 and alpha1 integrin I-domain in ELISA and FC studies.

**Antibody First Published in:** Hughes et al. Platelet integrin alpha2 I-domain specific antibodies produced via domain specific DNA vaccination combined with variable gene phage display. *Thromb Haemost.* 2005 Dec;94(6):1318-26. [PMID:16411412](#)

**Note on publication:** Describes the generation of mice scFvs by phage display and rabbit polyclonal antibodies raised against alpha2 integrin I-domain. The scFv specificity and reactivity was determined by ELISA and FC.

## Product Form

**Size:** 200 µg Purified antibody.

**Purification:** Protein A affinity purified

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.