



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-EDAR [EDAR12] Standard Size Ab00785-1.1

Isotype and Format: Mouse IgG1, Kappa

Clone Number: EDAR12

Alternative Name(s) of Target: ectodysplasin A receptor; Tumor necrosis factor receptor superfamily member EDAR; Anhidrotic ectodysplasin receptor 1; Downless; Ectodermal dysplasia receptor; Ectodysplasin-A receptor

UniProt Accession Number of Target Protein: Q9R187

Published Application(s): Agonist, SPR, WB, ELISA

Published Species Reactivity: Chicken, Dog, Rat, Human, Mouse

Immunogen: EDAR12 was prepared by immunizing female OVE1B mice (with the Edar gene deleted) subcutaneously with mouse EDAR-Fc and positive hybridoma clones were screened for binding mEDAR by ELISA.

Specificity: EDAR12 recognises and binds to CRD1 of the extracellular domain of mouse EDAR. EDAR3 cross-reacts with EDAR derived from human, dog, rat and chicken when EDAR is fused to the glycosylphosphatidylinositol anchor of TRAILR3. EDAR is the receptor for the TNF family ligand EDA1, which is a type II transmembrane protein possessing a collagen-like domain and a C-terminal TNF-homology domain. EDAR is important for the correct development of skin appendages including hair, teeth and eccrine sweat glands. LoF mutations in the Eda gene is known to cause XLHED (X-linked hypohidrotic ectodermal dysplasia), and results in abnormal development.

Application Notes: EDAR12 can be used in surrogate reporter assay in which Fas-sensitive cells were transfected with a construct formed by fusing the ectodomain of mouse or human EDAR to the intracellular domain of Fas. Binding of EDAR12 resulted in the induction of apoptosis in these cells, confirming the agonistic activity of the antibody. (EC50 (dose required to kill half of EDAR:Fas-expressing Jurkat cells) hEDAR:Fas ~ 5 ng/ml; EC50 mEDAR:Fas ~ 10 ng/ml - both were less active than DEAI-Fc). Tail-hair formation was also rescued after administration of EDAR3 to newborn EDA-deficient Tabby mice, with 0.35 mg/kg required for half-maximal tail-hair reversion. EDAR123 can also be used in ELISA and as a Fab fragment (generated by ficin digestion) in SPR experiments. EDAR can also be recognized by EDAR12 under both reducing (+DTT) and non-reducing conditions.

Antibody First Published in: Kowalczyk et al. Molecular and therapeutic characterization of anti-ectodysplasin A receptor (EDAR) agonist monoclonal antibodies. J Biol Chem. 2011 Sep 2;286(35):30769-79. [PMID:](#)

Note on publication: Describes the generation and characterization of anti-EDAR antibodies that mimic the action of EDA1 in stimulating EDAR both in vitro and in vivo.

Product Form

Size: 200 µg Purified antibody.

Purification:

Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.