



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Lymphotoxin alpha [3F12.2D3] Standard Size Ab00788-10.3

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This chimeric human antibody was made using the variable domain sequences of the original Mouse IgG2b format, for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Human IgG1, Fc Silent™, Kappa

Clone Number: 3F12.2D3

Alternative Name(s) of Target: chimeric 3F12;LT; LTα3; LTα; Tumor necrosis factor beta; Tumor necrosis factor β; Tumor necrosis factor-beta; tumor necrosis factor-β; TNF beta; TNF-beta; TNF β; TNF-β; Lymphotoxin-alpha; Lymphotoxin alpha

UniProt Accession Number of Target Protein: P01374

Published Application(s): FC;SPR, neutralize, ELISA

Published Species Reactivity: Human

Immunogen: 3F12 was prepared by hyperimmunizing BALB/c mice with purified recombinant human LTα expressed in E. coli - the hybridoma clone was selected for its ability to bind LTα3 by ELISA. The CDRs of the antibody were grafted to the framework regions of the mice mAb 2G7 to create the chimeric version of the antibody.

Specificity: 3F12 binds to human LTα3 and LTαβ (with a KD of ~0.3 nM, determined by BIACORE; ~ 37 pM, determined by ELISA) in both its soluble homotrimeric and membrane heterotrimeric forms. LTα belongs to the tumor necrosis factor (TNF) superfamily and is secreted as a homo-trimer (LTα3), or is expressed on the cell surface in complex with LTβ. Lymphotoxin is produced by lymphocytes and in its homotrimeric form binds to TNFRSF1A/TNFR1, TNFRSF1B/TNFB2 and TNFRSF14/HVEM. LTα3 signaling induces target cells to upregulate many chemokines and cytokines in an NFκB-dependent manner.

Application Notes: The properties of the chimeric 3F12 is equivalent to that of the original 3F12 mAb. 3F12 is able to bind to Th2 cells and additionally is can functionally block human LTα3 versus an isotype control (trastuzumab) - LTα3-induced ICAM upregulation is blocked on HUVEC (Human Umbilical Endothelial Cells) (IC50 0.04 nM) and LTα3-induced IL-8 is blocked in A549 cells. The antibody has also been used in ELISA studies to detect LT-alpha and to measure binding.

Antibody First Published in: Catellana et al. Resurrection of a Clinical Antibody: Template

ProteoGenomic de novoProteomic Sequencing and Reverse Engineering of an Anti-Lymphotoxin Alpha Antibody Proteomics. 2011 Feb; 11(3): 395–405. [PMID:21268269](#)

Note on publication: Describes the generation of anti-LT antibody and compares binding specificity and affinity of the chimeric antibody against the original antibody by BIACORE and ELISA.

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.