



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-MSP2 [6D8] Standard Size Ab00804-1.1

Isotype and Format: Mouse IgG1, Kappa

Clone Number: 6D8

Alternative Name(s) of Target: Merozoite surface protein 2

UniProt Accession Number of Target Protein: Q1WG90

Published Application(s): WB, ELISA, IF

Published Species Reactivity: Plasmodium falciparum

Immunogen: The 6D8 hybridoma cell line was generated from the fusion of Sp2/0 myeloma cells with spleen cells derived from female CBA mice immunized with polymeric recombinant 3D7 MSP2 (Ag1642). The hybridoma was selected for its ability to bind monomeric and polymeric Ag1642 (3D7 & FC27) by ELISA.

Specificity: 6D8 reacts with both 3D7 and FC27 recombinant MSP2, and preferentially with monomeric MSP2. The antibody binds to an epitope found at the conserved N-terminal region consisting of the sequence FINNAYNMSIRRS (aa 11-23) - the binding affinity for this sequence is 6 nM. MSP2 is a highly expressed and intrinsically disordered GPI-anchored membrane protein which is categorized into 2 allelic families/polymorphisms: 3D7 and FC27.

Application Notes: 6D8 reacts strongly and equivalently with both 3D7 and FC27 recombinant MSP2 by ELISA. 6D8 gives weak and no IFA signal on the parasite (i.e. not recombinant) FC27 and 3D7 MSP2 respectively, due to its inaccessibility on the merozoite surface. 6D8 can be used to detect recombinant FC27 and 3D7 MSP2 which has been subjected to SDS-PAGE under non-reducing conditions. Parasite FC27 MSP reactivity with 6D8 is enhanced in western blots when carried out in reducing conditions. 6D8 can be used to detect MSP2 by immunofluorescence.

Antibody First Published in: Adda et al. Antigenic characterization of an intrinsically unstructured protein, Plasmodium falciparum merozoite surface protein 2. Infect Immun. 2012 Dec;80(12):4177-85.

[PMID:22966050](#)

Note on publication: Describes the generation of a panel of antibodies against MSP2. Binding analyses of both recombinant and native MSP2 was performed.

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.