



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Podoplanin (MAP tag) [PMab-1] Bulk Size Ab00875-24.2-BT

This chimeric goat antibody was made using the variable domain sequences of the original rat IgG2a format, for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Goat IgG, Kappa

Clone Number: PMab-1

Alternative Name(s) of Target: PDPN; Aggrus; Glycoprotein 38; Gp38; OTS-8; Ots8; PA2.26 antigen; T1-alpha; T1A; Transmembrane glycoprotein E11

UniProt Accession Number of Target Protein: Q62011

Published Application(s): epitope tag, FACS, WB, Block, ELISA, FC, IHC

Published Species Reactivity: Mouse

Immunogen: This antibody was raised by immunising rats with 14-residue synthetic peptide mpp3851, which corresponds to amino acids 38-51 (GDGMVPPGIEDKIT) of the platelet-aggregation-stimulating (PLAG) domain of mouse podoplanin. Spleen cells were then harvested, and fused to P3U1 cells to generate stable hybridomas.

Specificity: This antibody is specific for amino acids 38-51 (GDGMVPPGIEDKIT) of the platelet-aggregation-stimulating (PLAG) domain of mouse podoplanin (PDPN). This antibody, therefore, possesses high affinity and specificity for the MAP epitope tag (GDGMVPPGIEDK), which is derived from the PLAG domain of murine podoplanin. PDPN is a type I transmembrane protein that interacts with the platelet receptor C-type lectin-like receptor-2 (CLEC-2). PDPN is expressed by a wide variety of cells, including podocytes, lymphatic endothelial cells and type I alveolar cells. PDPN is also upregulated on certain cancer cells, where it mediates tumour-induced platelet aggregation by binding to CLEC-2 on platelets. Chihara (2018) associated Podoplanin with a co-inhibitory receptor in T-cell activation along with PD-1, TIM-3, LAG-3 and TIGIT.

Application Notes: This antibody binds to the PLAG domain of murine PDPN and is, therefore, able to block its interaction with C-type lectin-like receptor-2 (CLEC-2) (Yamada, 2017). This has a number of potential therapeutic applications, including the suppression of lymphangiogenesis and inflammatory macrophage infiltration at the site of wound healing in mouse corneal suture and ear section models (Maruyama et al, 2014), as well as suppression of transplant rejection. This antibody possesses high affinity and specificity for the MAP epitope tag (GDGMVPPGIEDK), and can, therefore, be used to detect and purify tagged proteins in a range of assay formats (Fujii et al, 2016). PMab-1 performs well in immunohistochemical analysis, and has been used to stain the mouse melanoma B16-F10 and mouse

podoplanin-expressing CHO transfectants (Kaji et al, 2012). The use of this antibody in immunostaining of somatic tissues, including mouse lung lymphatic vessels, kidney glomeruli, pulmonary alveoli, pulmonary pleura and salivary gland myoepithelial cells, has been successfully performed on both paraffin-embedded sections and frozen sections (Kaji et al, 2012). The specificity of this antibody has been confirmed in Western blot analysis (Kaji et al, 2012), while Western blot analysis has also revealed that activation of LPS-induced inflammatory pathways, including the MAPK and NF- κ B pathways, is affected by PMab-1 (Maruyama et al, 2014). The mouse-rat chimeric antibody mPMab-1 is derived from PMab-1 (Yamada et al, 2017).

Antibody First Published in: Kaji et al Immunohistochemical Examination of Novel Rat Monoclonal Antibodies against Mouse and Human Podoplanin *Acta Histochem. Cytochem.* 45 (4): 227-237, 2012. doi:10.1267/ahc.12008 PMID:23012488

Note on publication: Describes the original generation and characterisation of this antibody.

Product Form

Size: 1 mg Purified antibody in bulk size.

Purification: Protein A affinity purified

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.