



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-IgG1 Fc [TP878] Standard Size Ab01437-10.159

This chimeric rabbit antibody was made using a variable domain sequence of the original VHH format, for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Human IgG1-Fc fusion

Clone Number: TP878

Alternative Name(s) of Target: immunoglobulin G1 fragment crystallizable region; immunoglobulin G1 Fc region; immunoglobulin G1

UniProt Accession Number of Target Protein:

Published Application(s): dot blot assay

Published Species Reactivity: Mouse

Immunogen: This antibody was raised by immunising alpacas with 1.0 mg polyclonal mouse IgG. Subsequently, the generation of nanobody immune libraries and the selection of antigen-specific nanobodies by phage display from these libraries were performed.

Specificity: This antibody recognises Fc fragment of the IgG1.

Application Notes: Anti-IgG nanobodies often perform more efficiently than polyclonal secondary antibodies in various experimental assays. Therefore, they are recommended for various experimental procedures. Anti-IgG nanobodies were shown to be more sensitive during Western blots when conjugated with different molecules (HRP, fluorophores etc.) (Pleiner et al., 2018). Furthermore, combination of two distinct nanobodies for immunofluorescence labelling provided exceptionally clean and strong signal, unprecedented among monovalent secondary antibodies (Pleiner et al., 2018). Additionally, using such an antibody allows skipping two-step incubation procedures during immunostaining as the nanobody can be first incubated with a primary antibody and this conjugate might be added to the cells in a one-step procedure (Pleiner et al., 2018). Finally, such nanobodies have been recommended as very effective for colocalisation studies when more than one fluorophore is needed (Pleiner et al., 2018).

Antibody First Published in: Pleiner et al. A toolbox of anti-mouse and anti-rabbit IgG secondary nanobodies. *J Cell Biol.* 2018 Mar 5;217(3):1143-1154. doi: 10.1083/jcb.201709115. Epub 2017 Dec 20.

[PMID:29263082](#)

Note on publication: The article describes the generation and characterisation of this nanobody.

Product Form

Size:

200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.