



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-CCR2 [AbAb-CCR2] Standard Size Ab02444-2.0

This chimeric mouse antibody was made using the variable domain sequences of the original rat IgG2b format, for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Mouse IgG2a, Kappa

Clone Number: AbAb-CCR2

Alternative Name(s) of Target: CD192; C-C chemokine receptor type 2; C-C CKR-2; CC-CKR-2; CCR-2; Monocyte chemoattractant protein 1 receptor; MCP-1-R; JE/FIC receptor; CMKBR2

UniProt Accession Number of Target Protein: P51683

Published Application(s): Block, FC

Published Species Reactivity: Mouse

Immunogen: The original antibody was generated by immunizing rats with CHO cells stably transfected with mouse CCR2.

Specificity: This antibody specifically binds mouse CCR2 and does not cross react with mouse CCR5 or human CCR2 and CCR5. CCR2 is a key functional receptor for CCL2 but can also bind CCL7 and CCL12 chemokines. Its binding with CCL2 on monocytes and macrophages mediates chemotaxis and migration. It regulates the expression of T-cell inflammatory cytokines and T-cell differentiation, promoting the differentiation of T-cells into T-helper 17 cells (Th17) during inflammation.

Application Notes: This antibody can be used to detect mouse CCR2 expressed on the surface of CHO cells and mouse peripheral blood leukocytes using flow cytometry. This antibody is capable of reducing the activity of murine monocyte chemotactic protein 1 in a manner similar to the well characterized anti-CCR2 antibody MC-21. This antibody can prevent the influx of monocytes in thioglycolate-induced peritonitis. This antibody is also capable of blocking the binding of mouse CCR2 with its ligand in vitro and in vivo. The blockade of CCR2 with this antibody also impairs monocytes/macrophage recruitment in vivo. This antibody can be used to determine the role of chemokine receptor CCR2 during initiation and progression of collagen induced arthritis in mice using blocking antibodies. This antibody is capable of abolishing long-term behavioral recovery after ischemic stroke and drastically decreased tissue expression of anti-inflammatory genes, including TGFβ, CD163, and Ym1.

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.