



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Thomsen-nouveaupolysaccharide A1 [KT-IgM-8] Standard Size Ab02511-10.0

Isotype and Format: Human IgG1, Kappa

Clone Number: KT-IgM-8

Alternative Name(s) of Target: Tn-PS A1; Thomsen-nouveau (Tn); Tn antigen; Tn saccharide; N-acetyl Galactosamine; Alpha-GalNAc; Alpha GalpNAc; GalNAc; GalNAc-alpha; N-acetyl-alpha-D-galactosamine; Blood group A antigen

UniProt Accession Number of Target Protein:

Published Application(s): Complement-dependent cytotoxicity, glycan array technologies, in vitro, in vivo, ELISA, FC

Published Species Reactivity: Human

Immunogen: The original antibody was generated by immunizing C57BL/6 mice against Tn-PS A1.

Specificity: The antibody is specific for Tn-PS A1. It does not cross react with PS A1.

Application Notes: The specificity of the original format of the antibody was confirmed by ELISA analysis. The ability of the antibody to bind cancer cells expressing the Tn antigen on the cell surface (MCF-7 and HCT-116) was evaluated by by FC. The antibody demonstrated the ability to bind both MCF-7 and HCT-116 tumor cells at 30 µg/mL with a shift in fluorescence of 49% in both cell lines. The antibody demonstrated in vitro tumor killing using an LDH assay. The assay measured that the antibody-induced complement-dependent cytotoxicity was 30%. In vivo passive immunotherapy approach using an MCF-7 cell line-derived xenograft model was evaluated. The antibody demonstrated approximately 40% reduction in tumor growth in the MCF-7 model (Trabbic et al., 2018; PMID: 30030557).

Antibody First Published in: Trabbic et al. Production of a mouse monoclonal IgM antibody that targets the carbohydrate Thomsen-nouveau cancer antigen resulting in in vivo and in vitro tumor killing Cancer Immunol Immunother. 2018 Sep;67(9):1437-1447. [PMID:30030557](#)

Note on publication: The original paper describes the generation and characterization of the antibody.

Product Form

Size: 100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.