



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-CD8 (Lyt-2.2) [2.43] Standard Size Ab02554-2.0

This chimeric mouse antibody was made using the variable domain sequences of the original Rat IgG2b format, for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Mouse IgG2a, Kappa

Clone Number: 2.43

Alternative Name(s) of Target: Lyt 2.2; T-cell surface glycoprotein CD8 alpha chain; Cd8a; T-cell surface glycoprotein Lyt-2; lyt-2; lyt2; CD_antigen: CD8a; 2.43.1

UniProt Accession Number of Target Protein: P01731

Published Application(s): Blocking, IP, FC, IF

Published Species Reactivity: Mouse

Immunogen: This antibody was generated by immunizing a Lewis rat with i.v. 3 times at 2-wk intervals with 15×10^6 L3 cloned cytolytic T cells. Three days after the third immunization, the rat spleen cells were fused with either P3 or SP2/0 cells using polyethylene glycol.

Specificity: This antibody is specific for the lyt-2 protein of mice (it is specific to the Lyt2.2 variant which is expressed in the mouse strains BALB/c and C57BL/6 (B/6)). Lyt-2 is an integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class I molecule:peptide complex.

Application Notes: This antibody was used for an immunofluorescence experiment on mouse thymocytes and mouse spleen cells (Sarmiento et al, 1980; pmid:6159417). Immunoprecipitation was performed on NP-40 extract of 126I surface labeled L3 cells using this antibody (Sarmiento et al, 1980; pmid:6159417). This antibody was used to try and block the cytolytic activity from L3 against P-185. It only blocks the receptor-mediated but not lectin mediated cytolysis by L3 (Dialynas et al, 1981; pmid:6788878). Flow cytometry was performed with this antibody on washed cells from the lymph node, spleen, thymus and peripheral blood from mice (Rodriguez and Sriram, 1988; pmid:2966203). This antibody was used to stain mouse spleen cells for use in flow cytometry (Mielke et al, 1988; pmid:2969373).

Antibody First Published in: Sarmiento et al. IgG or IgM monoclonal antibodies reactive with different determinants on the molecular complex bearing Lyt 2 antigen block T cell-mediated cytolysis in the absence of complement J Immunol. 1980 Dec;125(6):2665-72. [PMID:6159417](https://pubmed.ncbi.nlm.nih.gov/6159417/)

Note on publication: In the paper five rat monoclonal antibodies were derived that express specificities for determinants present on the molecular complex bearing the Lyt 2 antigen.

Product Form

Size: 100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration:

1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.