



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-CD56 ANK-1 [8G6] Standard Size Ab02588-23.0

Isotype and Format: Rabbit IgG, Kappa

Clone Number: 8G6

Alternative Name(s) of Target: CD56; ANK1; NCAM1; NCAM-1; N-CAM-1; Ncam; Neural cell adhesion molecule 1

UniProt Accession Number of Target Protein: P13595

Published Application(s): IHC (P), WB, FC

Published Species Reactivity: Human, Mouse

Immunogen: The original antibody was generated by immunizing rats with purified mouse brain N-CAM.

Specificity: This antibody binds an epitope called ANK-1 present on most NCAM isoforms expressed on a subset of NK cells of mouse and human. This epitope is selectively expressed on pre-activated, adherent natural killer NK cells (pre-A-NK cells), which are interleukin (IL)-2-stimulated NK cells, that are selectively endowed with the increased expression of integrins and ability to adhere to solid surfaces, migrate into, infiltrate, and destroy cancerous tissues.

Application Notes: This antibody was used to classify the nonactivated human peripheral blood NK cells expressing ANK-1 epitope using flow cytometry. It was reported that ANK-1 epitope was preferentially expressed by around 26% of NK cells (range 9–56%) and was exclusively CD56dim. This antibody was also capable of detecting ANK-1 epitope from crude human peripheral blood NK cell and affinity purified brain N-CAM lysates using western blot. This antibody was also used to test the ability of NK cells to respond to IL-2 by generation and expansion of A-NK cells with LAK activity, following depletion of ANK-1+ NK cells by treatment with mAb ANK-1 and complement. This antibody can also be used for the immunohistochemical staining of paraformaldehyde fixed paraffin embedded tissues of spleen, lymph nodes, tonsils, intestines, lungs, and liver tissues (PMID: 15356097).

Antibody First Published in: Shen Li et al. A novel epitope of N-CAM defines precursors of human adherent NK cells. *J Leukoc Biol.* 2004 Dec;76(6):1187-99. [PMID:15356097](#)

Note on publication: This paper studies a subpopulation of NK cells expressing a unique epitope ANK-1 using an antibody.

Product Form

Size: 100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In:

PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.