



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-CD227 [175C5] Standard Size Ab02886-1.1

Isotype and Format: Mouse IgG1, Kappa

Clone Number: 175C5

Alternative Name(s) of Target: Mucin-1; MUC1; Breast carcinoma-associated antigen DF3; Cancer antigen 15-3; CA 15-3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated epithelial membrane antigen; EMA; Tumor-associated mucin

UniProt Accession Number of Target Protein: P15941

Published Application(s): WB, FC, IHC

Published Species Reactivity: Human

Immunogen: The mouse version of this antibody was raised by immunizing a BALB/c mouse with milkfat globule membranes.

Specificity: This antibody is specific for human CD227. The alpha subunit of CD227 has cell adhesive properties. It can act both as an adhesion and an anti-adhesion protein. May provide a protective layer on epithelial cells against bacterial and enzyme attack.

Application Notes: To characterize 13 different ovarian cancer xenografts, immunohistochemistry was performed different human ovarian cancer cell lines using the mouse version of this antibody. It only stained most tissues moderately (Molthoff et al, 1991; pmid:1985883). To determine whether the appearance of tumour-associated macin epitopes can be explained by altered post translational modification of mucin in tumours, flow cytometry was used on EBY-immortalized B-cells and Burkitt's lymphomas using the mouse version of this antibody (Jerome et al, 1992; pmid:1382849). To study three antigenic systems for their use as tumour toxins, a western blot was performed on the human mammary carcinoma cell line MBMDA-23 1 and the colon carcinoma cell line SW 1222 using the mouse version of this antibody (Wick and Groner, 1997; pmid:9459206).

Antibody First Published in: Molthoff et al. Human ovarian cancer xenografts in nude mice: characterization and analysis of antigen expression Int J Cancer. 1991 Jan 2;47(1):72-9. [PMID:1985883](#)

Note on publication: 13 different human ovarian cancer xenografts grown subcutaneously in nude mice were characterized. The tumor lines represented 5 histological subtypes: serous, mucinous, clear-cell carcinoma, carcinosarcoma and undifferentiated.

Product Form

Size:

100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.