



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-15-acetyl-deoxynivalenol [NAT-267] Standard Size Ab02929-23.159

This chimeric rabbit antibody was made using the variable domain sequences of the original VHH format for improved compatibility with existing reagents assays and techniques.

Isotype and Format: Rabbit IgG-Fc fusion

Clone Number: NAT-267

Alternative Name(s) of Target: 15-AcDON; DON; Mycotoxin; Deoxynivalenol; Vomitoxin, Fusarium mycotoxin; mNAT-267; NAT-267; ChEBI: 10022

UniProt Accession Number of Target Protein:

Published Application(s): SPR, ELISA

Published Species Reactivity: Fusarium species

Immunogen: The original antibody was generated by immunization of a llama (*Llama glama*) with the protein conjugate 15-DON-BSA in a TiterMax Classic adjuvant.

Specificity: This antibody is specific for mycotoxin 15-acetyldeoxynivalenol (15-AcDON). 15-Acetyldeoxynivalenol belongs to the class of type B trichothecene mycotoxins, generated by several *Fusarium* fungi. Deoxynivalenol (DON), along with 3-acetyl-deoxynivalenol (3-ADON) and 15-acetyldeoxynivalenol (15-ADON), occur in grains and cereal products and is often hazardous to humans and livestock.

Application Notes: This antibody can bind 15-AcDON in a competitive direct enzyme-linked immunosorbent assay (CD-ELISA) and fluorescence polarization assay (FPA). This antibody does not cross react with DON; 3-acetyldeoxynivalenol (3-AcDON); neosolaniol (NEO); diacetoxyscirpenol (DAS); and T-2 toxin. Surface plasmon resonance (SPR) analysis of soluble NAT-267 binding kinetics to immobilized 15-DON-Horse Radish Peroxidase (HRP) indicated a dissociation constant K_D of 5µM (PMID: 18632156).

Antibody First Published in: Doyle et al. Cloning, expression, and characterization of a single-domain antibody fragment with affinity for 15-acetyl-deoxynivalenol. *Mol Immunol.* 2008 Aug;45(14):3703-13.

[PMID:18632156](#)

Note on publication: Describes the generation of a llama single-domain antibody fragment specific to the mycotoxin 15-acetyldeoxynivalenol.

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.