



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-IL-2 [UFKA-20] Bulk Size Ab02947-10.3-BT

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

Isotype and Format: Human IgG1, Fc Silent™, Kappa

Clone Number: UFKA-20

Alternative Name(s) of Target: IL2; IL-2; TCGF; Interleukin-2; T-cell growth factor; Aldesleukin

UniProt Accession Number of Target Protein: P60568

Published Application(s): ELISA, FC

Published Species Reactivity: Human

Immunogen: The original antibody was generated immunizing Balb/c mice with human IL-2.

Specificity: This antibody binds human IL-2. In particular, the antibody binds IL-2 dorsolaterally, with an angle of 55° counterclockwise to the vertical axis, contacting the C- and B-helices of IL-2 and thereby forming a “clamp” around the C-helix. The antibody engages IL-2 residues D84, N88, V91, E61, E68, and K64.

Application Notes: This antibody binds to Interleukin-2, a protein produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/UFKA-20 complex shows an improved selectivity towards regulatory T cells compared to IL-2 alone. IL-2/UFKA-20 can recognize the receptor IL of regulatory T cells on the base of the presence of CD25, therefore called CD25 bias. IL-2/UFKA-20 complex is able to efficiently stimulate CD25+ Treg cells in mice and rhesus macaques in vivo as well as in humans ex vivo.

Antibody First Published in: Karakus et al. Receptor-gated IL-2 delivery by an anti-human IL-2 antibody activates regulatory T cells in three different species. *Sci Transl Med.* 2020 Dec 16;12(574):eabb9283.

[PMID:33328333](#)

Note on publication: The authors identified a anti-human IL-2 antibody which was particularly efficient in delivering IL-2 to cells expressing trimeric IL-2Rs. Further IL-2/UFKA-20 complex, crystal structure, mechanism of action are described.

Product Form

Size: 1 mg Purified antibody in bulk size.

Purification: Protein A affinity purified

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer

storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.