



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Anti-PI-LPS [1E4] Standard Size Ab03230-23.0

This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG2a format for improved compatibility with existing reagents assays and techniques.

Isotype and Format: Rabbit IgG, Kappa

Clone Number: 1E4

Alternative Name(s) of Target: phase I lipopolysaccharide; LPS phase I; C. burnetii PI-LPS; C. burnetii phase I lipopolysaccharide

UniProt Accession Number of Target Protein:

Published Application(s): in vitro, in vivo, WB, ELISA

Published Species Reactivity: C. burnetii

Immunogen: The original antibody was generated by immunizing BALB/c mice with formalin-inactivated C. burnetii Nine Mile PI antigen.

Specificity: The antibody recognizes a PI specific epitope on PI-LPS. The antibody does not cross react with PII-LPS.

Application Notes: The specificity of the antibody (IgG2a) for PI-LPS was confirmed by ELISA analysis. The antibody detected PI-LPS by western blot analysis. The antibody was able to inhibit C. burnetii infection in vivo in a dose-dependent manner (Peng et al., 2012; PMID: 23053512). The original antibody could confer protection against C. burnetii aerosol infection. The Fab fragment, the scFv fragment and the humanized version of the antibody were constructed and characterized by indirect ELISA and Western Blot. They were able to bind to C. burnetii and to inhibit the infection in mice and in mouse Bone Marrow-Derived Macrophages (BMDM) in vitro. The humanized version inhibited C. burnetii infection in human macrophages in vitro. The Fab fragment was able to neutralize virulent C. burnetii resulting in inhibiting C. burnetii infection in both in vitro and in vivo systems (US20150087807A1). The ability of the original antibody, Fab, scFv, and humanized version of the antibody to inhibit C. burnetii infection in vivo was evaluated by comparing splenomegaly, bacterial burden, and pathological changes in the spleen with control mice at 14 days postinfection. IFA was used to analyze the ability of the original antibody to bind live virulent C. burnetii. The ability of the original to inhibit C. burnetii was higher than the ability of the other fragments. The humanized version of the antibody showed potential as a therapeutic against C. burnetii exposure (Peng et al., 2014; PMID: 25114119).

Antibody First Published in: Peng et al. Development of a lipopolysaccharide targeted peptide mimic vaccine against Q fever. J Immunol. 2012 Nov 15; 189(10): 10.4049/jimmunol.1201622.

[PMID:23053512](#)

Note on publication: The paper describes the generation and characterization of the antibody.

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.