



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-IL-2 [NARA1] Standard Size Ab03232-23.0

This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG2a format for improved compatibility with existing reagents assays and techniques.

Isotype and Format: Rabbit IgG, Kappa

Clone Number: NARA1

Alternative Name(s) of Target: IL2; IL-2; TCGF; Interleukin-2; T-cell growth factor; Aldesleukin

UniProt Accession Number of Target Protein: P60568

Published Application(s): crystallography, in vitro, in vivo, SPR, FC

Published Species Reactivity: Human

Immunogen: The original antibody was generated immunizing Balb/c mice with human IL-2.

Specificity: The antibody is specific for human IL-2. The antibody binds to a three-dimensional epitope of hIL-2 (residues P34, K35, R38, T41, Q74, and K76). The antibody overlaps with the CD25-binding epitope of IL-2. The antibody does not cross reacts with mouse IL-2.

Application Notes: SPR showed the binding of the antibody to IL-2 prevented further binding of hCD25 to IL-2; the affinity of the antibody to hIL-2 was $\sim 10^{-9}$ M. The crystal structure of the Fab fragment in complex with IL-2 was determined. IL-2/antibody complexes preferentially expand CD8⁺ T and NK cells in vivo while disfavoring CD25⁺ Tregs in transplantable metastatic melanoma model. This difference was also confirmed by kinetic proliferation studies. The complex IL-2/antibody decreased the tumor growth in mice by increasing the CD8⁺ T cell responses. The IL-2/antibody complex immunotherapy was also efficacious in a spontaneous melanoma model in mice. The complex IL-2/antibody shows potential as a therapeutic agent as it achieved strong antitumor immune responses (Arenas-Ramirez et al., 2016; PMID: 27903862). In order to prevent in vivo dissociation of the complex IL-2/antibody, NARA1leukin was constructed by integrating IL-2 into the antigen-binding groove on the antibody. The NARA1leukin version showed a longer half-life in vivo and more expansion of CD8⁺ T and NK cells, with no association with CD25, leading to a stronger antitumor response in B16-F10 murine models (Sahin et al., 2020; PMID: 33353953). The use of hIL-2/NARA1 in combination with Ezh2, an inhibitor of the histone methyltransferase enhancer of zeste homolog 2, showed enhanced tumor control in several mouse models of melanoma (Zingg et al., 2017; PMID: 28746871).

Antibody First Published in: Ramirez et al. Improved cancer immunotherapy by a CD25-mimobody conferring selectivity to human interleukin-2. *Sci Transl Med.* 2016 Nov 30;8(367):367ra166.

[PMID:27903862](#)

Note on publication: The paper describes the generation and characterization of the antibody, which is specific for IL-2 and it overlaps with the CD25-binding epitope of IL-2.

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.