



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-Beta Dystroglycan [AS30SS\_Hu6] Bulk Size Ab03277-10.3-BT

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

**Isotype and Format:** Human IgG1, Fc Silent™, Kappa

**Clone Number:** AS30SS\_Hu6

**Alternative Name(s) of Target:** Dystroglycan 1; Dag1; Beta-DG;  $\beta$ -DAG; BETA DAG; Dystroglycan; Dystrophin-associated glycoprotein 1; B-Dystroglycan; B Dystroglycan; Beta-dystroglycan;  $\beta$  dystroglycan;  $\beta$ -dystroglycan; AS30; humanized AS30

**UniProt Accession Number of Target Protein:** Q14118; Q62165

**Published Application(s):** IP, WB, ELISA, IF

**Published Species Reactivity:** Human, Mouse

**Immunogen:** The original antibody was generated by immunizing Balb/c and Trianni mice with human beta-dystroglycan. Humanized version was generated by grafting CDRs of parental mouse antibody AS30 onto human framework regions.

**Specificity:** This antibody recognizes human and mouse beta dystroglycan ( $\beta$ -DAG). The binding of this antibody to beta dystroglycan does not affect its binding to alpha dystroglycan. the Beta-dystroglycan is a transmembrane protein that plays an important role in connecting the extracellular matrix to the cytoskeleton. It plays an important role in linking dystrophin to the laminin-binding alpha-dystroglycan. It acts as a cell adhesion receptor in both muscle and non-muscle tissues. It also functions in cell adhesion-mediated signaling and implicated in cell polarity.

**Application Notes:** The binding specificity of this antibody for human and mouse beta dystroglycan (beta-DG) was confirmed using ELISA. This antibody is capable of immunoprecipitating both alpha-DG and beta-DG from C2C12 cell lysates, suggesting that the binding epitope of this antibody on beta dystroglycan is different from that of alpha dystroglycan. This antibody can also be used for the identification human beta-DG, mouse beta-DG and recombinant beta dystroglycan in a western blot. This antibody was also used for the immunofluorescent staining of unfixed human and mouse muscle tissues. This antibody was used in combination with anti-laminin-2 antibody C3\_Hu10 to generate a duobody called TBTI (US11254737).

**Antibody First Published in:** PMID:

**Note on publication:**

## Product Form

**Size:** 1 mg Purified antibody in bulk size.

**Purification:** Protein A affinity purified

**Supplied In:** PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.