

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



Anti-Cytokeratin 18 [RGE53] Standard Size, 100 µg, Ab03339-23.0 View online

Anti-Cytokeratin 18 [RGE53] Standard Size Ab03339-23.0

Isotype and Format: Rabbit IgG, Kappa

Clone Number: RGE53

Alternative Name(s) of Target: K18; CK-18; Keratin K18; Keratin type I cytoskeletal 18; Cell proliferationinducing gene 46 protein; Cytokeratin-18; Keratin-18

UniProt Accession Number of Target Protein: P05783

Published Application(s): WB, FC, IHC

Published Species Reactivity: canine, Chicken, Hamster, Rabbit, Rat, Swine, Zebrafish, Human, Mouse **Immunogen:** The original antibody was generated by immunizing a BALB/c mouse with a cytoskeletal preparation of HeLa cells.

Specificity: The antibody reacts exclusively with Cytokeratin 18 which is present in glandular epithelial cells of the digestive, respiratory, and urogenital tracts, endocrine and exocrine cells and mesothelial cells, as well as adenocarcinomas originating from them.

Application Notes: The antibody recognized glandular epithelial cells from female breast, digestive, respiratory, and urogenital tracts, endocrine and exocrine tissues, and mesothelial cells. Further, the antibody could distinguish adenocarcinomas and mesotheliomas, which stain positively in the indirect immunofluorescence technique (Ramaekers et al., 1983; PMID: 6193333). Further, the antibody produced fibrillar staining patterns in cultured HeLa cells, in cultured human hepatoma cells, F9 mouse teratocarcinoma cells and in a human colon carcinoma cell line (WiDr-218). The antibody detected Cytokeratin 18 in HeLa cells and cultured human hepatoma cells by western blot analysis (Ramaekers et al., 1983; PMID: 6350235). The antibody reacted with glandular and ductal epithelia, including those of the gastrointestinal tract, prostate, and kidney and with hepatocytes and mesothelium in immunohistochemistry (Ramaekers et al., 1990; PMID: 1690512).

Antibody First Published in: Ramaekers et al. Antibodies to intermediate filament proteins in the immunohistochemical identification of human tumors: an overview Histochem J. 1983 Jul;15(7):691-713. PMID:6350235

Note on publication: The original paper describes the generation and characterization of the antibody.

Product Form

Size: 100 μg Purified antibody. **Purification:** Protein A affinity purified **Supplied In:** PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at - 20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.