

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Anti-HLA-DR [HD8] Bulk Size Ab03367-11.3-BT

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

Isotype and Format: Human IgG2, Fc Silent[™], Kappa

Clone Number: HD8

Alternative Name(s) of Target: CD74; Beta chain HLA-DR; HLA-DR β-chain; β-chain; HLA class II

histocompatibility antigen gamma chain; HLA-DR antigens-associated invariant chain; la antigen-associated

invariant chain; li; p33

UniProt Accession Number of Target Protein: P04233

Published Application(s): immunotherapy, WB, FC

Published Species Reactivity: Human

Immunogen: The original antibody was generated by immunizing a human transgenic KM mice with HLA-DR-expressing L929 cells.

Specificity: This antibody specifically binds an epitope comprising amino acids 61-WNSQKDILEQARA-73 in the beta chain of human HLA-DR. This antibody can recognize most polymorphic forms of the human HLA-DR. This antibody can recognize 99.2% of HLA-DRB alleles and also is capable of cross reacting with HLA-DP alleles some HLA-DQ alleles. HLA-DR is an MHC class II cell surface receptor encoded by the human leukocyte antigen complex on chromosome 6 region 6p21.31. HLA-DR is also involved in several autoimmune conditions, disease susceptibility and disease resistance. The primary function of HLA-DR is to present peptide antigens, potentially foreign in origin, to the immune system for the purpose of eliciting or suppressing T-(helper)-cell responses that eventually lead to the production of antibodies against the same peptide antigen.

Application Notes: The binding specificity of this antibody to HLA-DR-positive B-lymphoblastoid cell line, SKW6.4 was tested using western blot. The reactivity of this antibody towards various HLA-DR positive cell lines like ARH77, Daudi, Granta519, IM-9, MC/CAR, Raji, RPMI1788, HS-Sultan, Namalwa, RL, Ramos, SKW6.4 etc. was determined using Alexa488 labelled antibody by flow cytometry. This antibody is reported to exert strong antibody dependent cellular cytotoxicity and complement-dependent cytotoxicity in vitro, and significantly extended the life span of immunocompromised mice inoculated with non-Hodgkin lymphoma cell lines. This antibody has potential applications in HLA-DR targeted immunotherapy as it is likely to evoke similarly strong responses in individuals carrying different HLA-DR alleles (PMID: 17428256). The Pro331Ser and Lys322Ala modified version of this antibody was incapable of expressing CDC in vitro and did not induce severe infusion reactions in rats and monkeys, even at extremely high doses. This

antibody still retained its Ab-dependent cellular cytotoxicity function as well as its antitumor activity in a tumor-bearing mouse model (PMID: 18250438).

Antibody First Published in: Tawara et al. Fully human antibody exhibits pan-human leukocyte antigen-DR recognition and high in vitro/vivo efficacy against human leukocyte antigen-DR-positive lymphomas.

Cancer Sci. 2007 Jun;98(6):921-8. PMID:17428256

Note on publication: Describes the generation and characterization of this antibody.

Product Form

Size: 1 mg Purified antibody in bulk size. **Purification:** Protein A affinity purified

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommed this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.