



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-NKp46 [NKP46-4] Standard Size Ab03381-10.0

This full-length reformatted human antibody was made using the variable domain sequences of the original Human scFv format for improved compatibility with existing reagents assays and techniques.

**Isotype and Format:** Human IgG1, Kappa

**Clone Number:** NKP46-4

**Alternative Name(s) of Target:** Natural cytotoxicity triggering receptor 1; Lymphocyte antigen 94 homolog; NK cell-activating receptor; Natural killer cell p46-related protein; NK-p46; hNKp46; CD335

**UniProt Accession Number of Target Protein:** O76036

**Published Application(s):** ELISA, FC

**Published Species Reactivity:** rhesus macaque, Human, Cynomolgus Monkey

**Immunogen:** The original antibody was isolated using phage display technology after three rounds of panning against human NKp-46.

**Specificity:** The antibody is specific for human NKp46. The antibody cross-reacts with cynomolgus monkey and rhesus macaque NK46 but does not cross-react with mouse NKp46. NKp46 is a receptor expressed by NK cells.

**Application Notes:** The binding specificity of this antibody towards human, cynomolgus monkey and rhesus monkey NKp-46 was determined using ELISA and flow cytometry. The binding of this antibody to NKp-46 expressed on the surface of CHOK1 and NK cells was confirmed using flow cytometry. The original IgG antibody bound ectodomain of human NKp-46 with a binding affinity of  $K_d = 1.1$  nM. This binding of this antibody to NK cells resulted in an increase in the interferon gamma release (CN113637074).

**Antibody First Published in:** [PMID:](#)

**Note on publication:**

## Product Form

**Size:** 200 µg Purified antibody.

**Purification:** Protein A affinity purified

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic

procedures for humans or animals.