



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-Carcinoembryonic Antigen [COL-1] Bulk Size Ab03385-23.0-BT

This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG2a format for improved compatibility with existing reagents assays and techniques.

**Isotype and Format:** Rabbit IgG, Kappa

**Clone Number:** COL-1

**Alternative Name(s) of Target:** CEA

**UniProt Accession Number of Target Protein:** Q13984

**Published Application(s):** in vivo, RIA, therapeutic, WB, IF, IHC

**Published Species Reactivity:** Human

**Immunogen:** The original antibody was generated by immunizing BALB/c mice with extracts or membrane-enriched fractions of biopsy material from either primary or metastatic colon carcinoma lesions.

**Specificity:** The antibody is specific for CEA. COL-1 does not cross-react with NCA, NFA-1, BGP and PMNs. CEA is one of the most commonly used tumor markers in serum immunoassay determinations of carcinoma.

**Application Notes:** The antibody reacted with purified carcinoembryonic antigen in solid-phase radioimmunoassay and by Western Blot analysis. The antibody was tested by fluorescence-activated cell sorting for cell surface binding to the LS174T WiDr and HT-29 colon carcinoma cell lines. The antibody was tested by FACS for binding to the surface of normal human polymorphonuclear leukocytes where it appeared nonreactive (Muraro et al., 1985; PMID: 2413997). The antibody was shown to bind CEA with a  $K_a$  of  $1.4 \times 10^9 \text{ M}^{-1}$  (Kuroki et al., 1989; PMID: 2474516). The antibody was used for immunohistochemistry on colorectal, gastric, and lung adenocarcinoma tumor tissues. In addition, pancreatic, breast, gallbladder, bile duct, malignant ovarian teratoma, mucinous endometrial carcinoma, colon cancer, remote colon mucosa and bladder cancer stained positive (Shi et al., 1994; PMID: 7520463 and Robbins et al., 1993; PMID: 8386136). The antibody reacted by immunofluorescence with a mouse colon adenocarcinoma MC-38 expressing high levels of cell-surface CEA cell line, but not with the NCA or BGP transfectants (Robbins et al., 1993; PMID: 8386136). 125I-labeled COL-1 IgG was shown to efficiently and specifically target the LS-174T human colon carcinoma xenograft in athymic mice. 131I-labeled COL-1 demonstrated a reduction of tumor growth rate in the same model (Siler et al., PMID: 8292968). 1131I-labeled COL-1 was tested in phase I trial in patients with gastrointestinal malignancies (Yu et al. 1996). The humanized version of the antibody was constructed and used in the detection and treatment of a CEA-expressing tumor (US8828717B2).

**Antibody First Published in:** Muraro et al. Definition by monoclonal antibodies of a repertoire of epitopes

on carcinoembryonic antigen differentially expressed in human colon carcinomas versus normal adult tissues Cancer Res. 1985 Nov;45(11 Pt 2):5769-80. [PMID:2413997](#)

**Note on publication:** The original paper describes the generation and characterization of a set of antibodies. The antibodies show selective reactivity for human colon carcinomas.

## Product Form

**Size:** 1 mg Purified antibody in bulk size.

**Purification:** Protein A affinity purified

**Supplied In:** PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.