



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-OPGbp [AT] Bulk size M Ab03389-10.29-BT

This is a Fab fragment with His tag.

Isotype and Format: Human Fab fragment, His-Tagged, Kappa

Clone Number: AT

Alternative Name(s) of Target: Osteoprotegerin binding protein; Tumor necrosis factor receptor superfamily member 11B; Osteoclastogenesis inhibitory factor; Osteoprotegerin

UniProt Accession Number of Target Protein: O00300

Published Application(s): inhibition, SPR, ELISA

Published Species Reactivity: Human

Immunogen: The original antibody was generated by panning against a human Fab library.

Specificity: The antibody recognizes Human OPGbp (residues 143-317). OPGbp acts as decoy receptor for TNFSF11/RANKL and thereby neutralizes its function in osteoclastogenesis. It inhibits the activation of osteoclasts and promotes osteoclast apoptosis in vitro.

Application Notes: The specificity of the original format of the antibody was confirmed by ELISA analysis. The binding affinity of the Fab and IgG1 fragments of the antibody to OPGbp was measured by surface plasmon resonance ($K_d \approx 140$ nM and 0.33-0.43 nM respectively). The interaction of human OPGbp and its receptor ODAR was measured in a fluorescence energy transfer (FRET) assay. The Fab fragment of the antibody inhibited OPGbp binding to soluble ODAR-Fc fusion with an IC_{50} of 550 nM. Bone Marrow Assay using the Fab fragment of the antibody gave a 50% decrease (IC_{50}) at 57.8 nM. In RAW Cell Assay the Fab fragment of the antibody showed an IC_{50} of 15 μ g/ml, 300 nM. In RAW Cell Assay, the 3 full length versions of the antibody "AT"405, "AT"406 and "AT"407 showed IC_{50} of 20.1 nM, 60.3 nM & 21.4 nM, respectively. "AT"405 or "AT"407 were tested in murine bone marrow assay, showing IC_{50} of 2.15 μ g/ml (14.4 nM), and 1.85 μ g/ml (12.5 nM), respectively. "AT"406 was tested in the bone marrow coculture assay and inhibited TRAP formation with an IC_{50} of 2.35 μ g/ml (14.4 nM) (US20140147444).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 1 mg Purified antibody in bulk size.

Purification: Purified by Immobilized Metal Affinity Chromatography

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without

added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: See vial label

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.