



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-EP-CAM [4B5E6D7] Standard Size Ab03393-1.1

**Isotype and Format:** Mouse IgG1, Kappa

**Clone Number:** 4B5E6D7

**Alternative Name(s) of Target:** CGMCC No.17406; Epithelial cell adhesion molecule; Adenocarcinoma-associated antigen; Cell surface glycoprotein Trop-1; Epithelial cell surface antigen; Epithelial glycoprotein; EGP; Epithelial glycoprotein 314; EGP314; hEGP314; KS 1/4 antigen; KSA; Major gastrointestinal tumor-associated protein GA733-2; Tumor-associated calcium signal transducer 1; CD326; EPCAM; MOC-31

**UniProt Accession Number of Target Protein:** P16422

**Published Application(s):** WB, ELISA, IHC

**Published Species Reactivity:** Human

**Immunogen:** The original antibody was generated by immunizing BALB/c mice with EP-CAM.

**Specificity:** The antibody is specific for EP-CAM. EP-CAM is a transmembrane glycoprotein mediating Ca<sup>2+</sup>-independent homotypic cell-cell adhesion in epithelia. EpCAM is also involved in cell signaling, migration, proliferation, and differentiation. It may act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. It plays a role in embryonic stem cells proliferation and differentiation and it up-regulates the expression of FABP5, MYC and cyclins A and E.

**Application Notes:** The antibody 4B5E6D7 and the commercial antibody MOC-31 were used to stain lung squamous cell carcinoma, lung adenocarcinoma, cervical cancer, colorectal carcinoma, duodenum and rectal tissues. Results showed the antibody specificity is comparable to the commercial antibody. Further, the sensitivity of the antibody 4B5E6D7 was higher than the commercial antibody. The antibody and the commercially available antibody MOC-31 were simultaneously detected on the normal tissue chip, indicating that the specificity of the antibody in normal tissue was equivalent to that of the commercially available antibody (CN110922485B).

**Antibody First Published in:** [PMID:](#)

**Note on publication:**

### Product Form

**Size:** 200 µg Purified antibody.

**Purification:** Protein A affinity purified

**Supplied In:**

PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.