

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



Anti-CD38 [13B5B11C8] Standard Size, 200 µg, Ab03398-23.0 View online

Anti-CD38 [13B5B11C8] Standard Size Ab03398-23.0

This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG1 format for improved compatibility with existing reagents assays and techniques.

Isotype and Format: Rabbit IgG, Kappa

Clone Number: 13B5B11C8

Alternative Name(s) of Target: CGMCC NO.22319; ADP-ribosyl cyclase/cyclic; 2'-phospho-ADP-ribosyl cyclase; 2'-phospho-ADP-ribosyl cyclase/2'-phospho-cyclic-ADP-ribose transferase; 2'-phospho-cyclic-ADP-ribose transferase; ADP-ribosyl cyclase 1; ADPRC 1; Cyclic ADP-ribose hydrolase 1; cADPr hydrolase 1; T10; 38C03

UniProt Accession Number of Target Protein: P28907

Published Application(s): WB, ELISA, IHC

Published Species Reactivity: Human

Immunogen: The original antibody was generated by immunizing BALB/c mice with CD38.

Specificity: The antibody is specific for CD38. CD38 is a single-chain transmembrane type II glycoprotein with a molecular weight of 45kD, including a long carboxyl-terminal extracellular domain (258 amino acids), a transmembrane domain (21 amino acids) and a short amino-terminal Intracellular region (21 amino acids). CD38 is a multifunctional protein with receptor and enzyme-mediated functions, which can act as receptor, adhesion molecule and exonuclease.

Application Notes: The antibody specifically detected CD38 by Western blot analysis. The antibody 13B5B11C8 and the commercial antibody 38C03 were used to stain B-cell lymphoma and T-cell lymphoma. Results showed the antibody specificity is comparable to the commercial antibody. Further, the sensitivity of the antibody 13B5B11C8 was higher than the commercial antibody. The antibody and the commercially available antibody 38C03 were simultaneously detected on the normal tissue chip, indicating that the specificity of the antibody in normal tissue was equivalent to that of the commercially available antibody (CN113912727A).

Antibody First Published in: PMID: Note on publication:

Product Form

Size: 200 µg Purified antibody. Purification:

Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at - 20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.