



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-CD40 [3B2] Standard Size Ab03420-23.0

This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG1 format for improved compatibility with existing reagents assays and techniques.

**Isotype and Format:** Rabbit IgG, Kappa

**Clone Number:** 3B2

**Alternative Name(s) of Target:** TNFRS5; Tumor necrosis factor receptor superfamily member 5; B-cell surface antigen CD40; Bp50; CD40L receptor CDw40; CD154R; 19-3B2

**UniProt Accession Number of Target Protein:** P25942

**Published Application(s):** therapeutic, FC

**Published Species Reactivity:** Primate, Human

**Immunogen:** The original antibody was generated by immunizing mice with recombinant human CD40 protein.

**Specificity:** This antibody recognizes and binds human and primate CD40, but does not cross react with mouse CD40. CD40 acts as a receptor for TNFSF5/CD40LG. This protein is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation.

**Application Notes:** The reactivity of this antibody to human, Ganlamp monkey and mouse CD40 protein expressed on the surface of CHO cells was determined using flow cytometry. This antibody positively binds human and primate CD40 proteins and does not cross react with mouse CD40. The original mouse IgG antibody binds human CD40 with a binding affinity of  $k_d = 6.13 \times 10^{-10}$  M. The tumor inhibitory activity of this antibody was also tested in vivo in B-HCD40 mice injected with murine MC-38 cells (colon cancer cells). This antibody exhibited good tumor inhibitory effect (CN111763259).

**Antibody First Published in:** [PMID:](#)

**Note on publication:**

### Product Form

**Size:** 200 µg Purified antibody.

**Purification:** Protein A affinity purified

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.