

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Anti-Apical complex glycoprotein CSL [3E2] Ab03423-1.1-BT

This reformatted mouse antibody was made using the variable domain sequences of the original Mouse IgM format for improved compatibility with existing reagents assays and techniques.

Isotype and Format: Mouse IgG1, Kappa

Clone Number: 3E2

Alternative Name(s) of Target: CSL; αCSL; Apical complex glycoprotein, CSL (circumsporozoite-like)

antigen

UniProt Accession Number of Target Protein:

Published Application(s): electron microscopy, therapeutic, WB, ELISA

Published Species Reactivity: Cryptosporidium parvum

Immunogen: The original antibody was generated by immunizing mice with GP25-200 of Cryptosporidium parvum.

Specificity: This antibody binds the apical complex glycoprotein CSL of Cryptosporidium parvum. The apicomplexan protozoan Cryptosporidium parvum is a coccidian parasite that infects intestinal epithelium and causes diarrheal disease in humans and other agriculturally important food animals for which specific therapy and immunoprophylaxis are unavailable. Cryptosporidiosis is self-limiting in hosts with normal immune systems; however, in neonates, the elderly, and hosts having congenital or acquired immunodeficiency diseases or chemotherapy-induced immunosuppression, cryptosporidial enterocolitis may become chronic and have severe consequences.

Application Notes: Apical complex glycoprotein antigen of Cryptosporidium parvum is critical to infectivity and targets protective immunity. This antibody exhibited morphologic changes in sporozoites and merozoites characterized by rapid and progressive formation, posterior movement, and release of membranous Ag-mAb precipitates. This antibody was capable of recognizing 46-770 kDa sporozoite antigens and 1300 kDa antigen soluble glycoprotein exoantigen (CSL) expressed by merozoites in a western blot (PMID: 9029117). This antibody elicited circumsporozoite precipitate (CSP)-like reaction which is characterized by the progressive posterior formation and release of membranous antigen-MAb complexes, after which zoites are rendered noninfective. This antibody is capable of neutralizing C. parvum in vitro and also capable of passively protecting against oocyst challenge and infection in vivo. The identification of sporozoite-neutralizing antibody was done using ELISA (PMID: 10768951). This antibody was reported to be highly effective in in reducing, but not eliminating, persistent C. parvum infection (PMID: 11796330). This antibody in combination with PLA2 or LL37 was also used in the generation of a biocide

fusion proteins to kill C. parvum sporozoites in vitro and in vivo (PMID: 20086143).

Antibody First Published in: Riggs et al. Protective monoclonal antibody defines a circumsporozoite-like glycoprotein exoantigen of Cryptosporidium parvum sporozoites and merozoites. J Immunol. 1997 Feb 15;158(4):1787-95. PMID:9029117

Note on publication: Describes the generation of neutralizing antibodies against C. parvum and their role in providing passive immunization against cryptosporidiosis.

Product Form

Size: 1 mg Purified antibody in bulk size. **Purification:** Protein A affinity purified

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommed this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.