



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Glypican-1 [AbAb-GPC1] Standard Size Ab03427-23.0

Isotype and Format: Rabbit IgG, Kappa

Clone Number: AbAb-GPC1

Alternative Name(s) of Target: GPC1

UniProt Accession Number of Target Protein: P35052

Published Application(s): ELISA, FC

Published Species Reactivity: Human

Immunogen: The original antibody was generated by immunizing 6-week-old healthy New Zealand female white rabbits subcutaneously with human glypican-1.

Specificity: This antibody binds human glypican-1, which is a cell surface proteoglycan that bears heparan sulfate. This protein is required for skeletal muscle differentiation. Glypican-1 (GPC1) was significantly expressed in serum exosomes of patients with pancreatic cancer and so GPC1 is considered as a tumor marker for pancreatic cancer.

Application Notes: The binding characterization of this antibody to human glypican-1 was determined using ELISA. The binding of this antibody to human glypican-1 expressed on the surface of human breast cancer cell line MDA-MB-231 was determined using flow cytometry (CN112300283).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.