



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-CXCL10 [2A] Ab03438-7.1-BT

This chimeric rat antibody was made using the variable domain sequences of the original Human IgG1 format for improved compatibility with existing reagents assays and techniques.

**Isotype and Format:** Rat IgG2a, Kappa

**Clone Number:** 2A

**Alternative Name(s) of Target:** IP-10; INP10; Mob1; SCYB10; Gamma-IP10; C-X-C motif chemokine 10; 10 kDa interferon gamma-induced protein; Interferon-inducible protein 10; Protein Mob-1; Small-inducible cytokine B10

**UniProt Accession Number of Target Protein:** P48973

**Published Application(s):** functional assay, therapeutic, ELISA

**Published Species Reactivity:** Rat

**Immunogen:** The original antibody was isolated from a human scFv phage library by panning against purified rat CXCL10. Later on the scFv was reformatted to whole IgG1.

**Specificity:** This antibody binds rat CXCL10. This pro-inflammatory cytokine is involved in a wide variety of processes such as chemotaxis, differentiation, and activation of peripheral immune cells, regulation of cell growth, apoptosis and modulation of angiostatic effects. This protein plays an important role during viral infections by stimulating the activation and migration of immune cells to the infected sites. Binding of CXCL10 to the CXCR3 receptor activates G protein-mediated signaling and results in downstream activation of phospholipase C-dependent pathway, an increase in intracellular calcium production and actin reorganization.

**Application Notes:** The binding affinity of this antibody to rat CXCL10 was confirmed using ELISA. The IgG version of this antibody binds rat CXCL10 with a binding affinity of  $K_d = 150$  pM. This antibody was capable of inhibiting migration of EL4 cells by CXCL-10 in a migration assay. Quantum dot nanocrystals labelled (Q-Dot-labeled) 2A was to study the movement and distribution of this antibody in collagen induced arthritis animal model using in vivo imaging technique. It was reported that the Q-Dot-labeled antibody was specifically accumulated in the joint area of the forefoot and lower foot, which are inflammation sites. The therapeutic effect of this antibody on arthritis was also studied in preclinical animal models of autoimmune arthritis. It was reported that there was a reduction in cell infiltration in a concentration dependent manner in the anti-CXCL-10-2A treated group, and the destruction of cartilage was also alleviated when compared to the IgG control treated control group. In the joints of the Anti-CXCL-10-2A injection group, the expression

of inflammatory cytokines IL-17, TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 was low (KR1020130121601).

**Antibody First Published in:** [PMID:](#)

**Note on publication:**

## Product Form

**Size:** 1 mg Purified antibody in bulk size.

**Purification:** Protein A affinity purified

**Supplied In:** PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.