



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-LAG-3 [ADI-31798] Ab03439-10.3-BT

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This reformatted human antibody was made using the variable domain sequences of the original Human IgG4 format for improved compatibility with existing reagents assays and techniques.

Isotype and Format: Human IgG1, Fc Silent™, Kappa

Clone Number: ADI-31798

Alternative Name(s) of Target: CD223; LAG3; FDC; Lymphocyte activation gene 3 protein

UniProt Accession Number of Target Protein: P18627

Published Application(s): ELISA, FC

Published Species Reactivity: Human

Immunogen: The original antibody was isolated from a yeast display library by two rounds of screening against human LAG-3.

Specificity: This antibody binds human LAG-3. The human LAG-3 is a cell surface molecule with diverse biologic effects on T cell function. It is an immune checkpoint receptor on activated T-cells.

Application Notes: This antibody can bind human LAG-3 in an ELISA. This antibody binds human LAG-3 with an affinity of $K_d = 9.73 \times 10^{-10}$ nM. The binding specificity of this antibody to human LAG-3 expressed on the surface of HEK293 and activated CD4+ T cells was done using flow cytometry. This antibody binds hLAG-3 expressed on the surface of HEK293 cells with an EC50 of 0.201 nM and activated CD4+ T cells with an EC50 of 0.00595 nM. This antibody can also effectively block the binding of LAG-3 and its ligand MHCII (HLA-DR) with an IC50 of 4.701 nM. The tumor suppressive effects of this antibody was studied in A375 human PBMC model (CN112166125).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 1 mg Purified antibody in bulk size.

Purification: Protein A affinity purified

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without

added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.