



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Glycoprotein IIb/IIIa [SE] Standard Size Ab03507-10.3

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This is a reformatted human IgG1 Fc Silent Fc Silent™ antibody, based on the original human scFv format, created for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Human IgG1, Fc Silent™, Lambda

Clone Number: SE

Alternative Name(s) of Target: GPIIb/IIIa; CD41/CD61; integrin α IIb β 3

UniProt Accession Number of Target Protein: P08514

Published Application(s): FC

Published Species Reactivity: Human

Immunogen: The original antibody was generated by screening human scFv antibody libraries.

Specificity: This antibody binds the glycoprotein GPIIb/IIIa complex (CD41/CD61 complex). The GPIIb/IIIa is a heterodimeric complex formed after synthesis of one β and one α subunit. The principal ligand for GPIIb/IIIa is fibrinogen, but it also binds to fibronectin, von Willebrand factor, vitronectin, thrombospondin and CD40 ligand. The binding between GPIIb/IIIa and fibrinogen dimers leads to platelet aggregation and thrombus formation, and this is possible only when the receptor adopts its activated conformation.

Application Notes: The antibody bound to activated platelets and inhibited platelet activation. This antibody was used for detection of activated platelets by flow cytometry (WO2020237304).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic

procedures for humans or animals.