

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



## Anti-S protein (RBD) [aRBD-7] Standard Size Ab03533-23.159

Isotype and Format: Rabbit IgG-Fc fusion

#### Clone Number: aRBD-7

**Alternative Name(s) of Target:** SARS CoV 2 S glycoprotein; COVID-19 Spike protein; RBD; Receptor Binding Domain; E2 glycoprotein; E2; Human coronavirus 2 spike glycoprotein; Peplomer protein; S glycoprotein; SARS coronavirus 2 S protein; SARS coronavirus 2 Spike Protein; SARS CoV 2 Spike protein; SARS CoV 2; SARS-CoV-2 S protein; SARSCoV2; SARS-COV-2 S protein; SARS-COV-2 Spike glycoprotein; SARSCOV2 Spike protein; Severe acute respiratory syndrome 2 spike glycoprotein; Severe acute respiratory syndrome virus 2 spike glycoprotein; Spike glycoprotein; 2019-nCoV; SARS-CoV2

UniProt Accession Number of Target Protein: P0DTC2

Published Application(s): Blocking, neutralizing, SPR, ELISA

### Published Species Reactivity: SARS-CoV-2

**Immunogen:** The original antibody was generated by immunizing two alpacas with highly purified recombinant SARS-CoV-2 RBD, and developing a VHH library. The antibody was isolated by panning against SARS-CoV-2 RBD.

**Specificity:** The antibody binds the Spike protein of the SARS-CoV-2.

Application Notes: The VHH with IgG1 Fc formats demonstrated strong binding to both RBD and the entire ectodomain (S1+S2) of SARS-CoV-2 spike in ELISA, with a low nanomolar 50% effective concentration (EC50 respectively 0.522 nM and 0.748 nM). The binding affinity of the VHH format and VHH-Fc formats to RBD were also measured using surface plasmon resonance, with KD values of 3.31 nM and 72 pM respectively. Both VHH and Fc fusion formats could block RBD-ACE2 interaction in a dose-dependent manner, as characterized by competitive ELISA. Furthermore, the fusion of two VHHs with nonoverlapping epitopes resulted in the hetero-bivalent VHHs, aRBD-2-7, which showed significantly higher RBD binding affinities (KD of 0.25 nM). Similarly, their Fc fusions also showed enhanced binding affinities, with a KD value of 0.22 nM. The homo- and hetero-bivalent VHHs exhibited potent neutralizing ability against SARSCoV-2 inoculated onto Vero E6 cells. The 50% neutralization dose was 3.18 ng/ml (0.111 nM), for aRBD-2-7. Finally, the ND50s for aRBD-2-7.Fc was 6.76 ng/ml (0.0606 nM) (Ma et al, 2021; PMID:33658349).
Antibody First Published in: Ma et al. Potent Neutralization of SARS-CoV-2 by Hetero-bivalent Alpaca Nanobodies Targeting the Spike Receptor-Binding Domain J Virol. 2021 Mar 3;95(10):e02438-20.
PMID:33658349

**Note on publication:** The original paper describes the generation and characterization of the antibody

### **Product Form**

**Size:** 100 μg Purified antibody. **Purification:** Protein A affinity purified

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at - 20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.