



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Hepcidin-25 [OH4] Ab03566-23.0-BT

This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG1 format, for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Rabbit IgG, Kappa

Clone Number: OH4

Alternative Name(s) of Target: HAMP; hHepc; HEPC; LEAP1; LEAP-1; Hepcidin; Liver-expressed antimicrobial peptide 1

UniProt Accession Number of Target Protein: P81172

Published Application(s): ELISA

Published Species Reactivity: Human

Immunogen: The original antibody was generated by immunizing mice with immunized with a N-terminal hepcidin peptide 'DTHFPIC' conjugated with a five amino acid peptide linker (i.e., GP GPG) to an OVA peptide (amino acids 323-336) sequence, i.e., full-length immunogen DTHFPICGP GPGISQAVHAAHAEINE.

Specificity: This antibody binds human hepcidin-25 near the N-terminus region and does not cross react with human hepcidin-20 or human hepcidin-22. This antibody also does not cross react with mouse or rat hepcidin-25. Hepcidin is a an iron-regulating peptide hormone made in the liver. Hepcidin is a cationic, cysteine-rich and tightly folded peptide stabilized by 4 disulfide bonds that plays a major role in innate immunity and iron homeostasis. It controls the delivery of iron to blood plasma from intestinal cells absorbing iron, from erythrocyte-recycling macrophages, and from iron-storing hepatocytes. Hepcidin acts by binding to and inactivating the sole cellular iron exporter, ferroprotein, which delivers iron to plasma from all iron-transporting cells.

Application Notes: The original mouse IgG1 antibody binds human hepcidin-25 with a binding affinity of $k_d = 1.2$ nM. The binding characterization of this antibody towards human hepcidin was done using ELISA (US8609817).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 1 mg Purified antibody in bulk size.

Purification: Protein A affinity purified

Supplied In:

PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.