



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-Hepcidin-25 [AN-LP1] Ab03568-1.1-BT

**Isotype and Format:** Mouse IgG1, Kappa

**Clone Number:** AN-LP1

**Alternative Name(s) of Target:** HAMP; hHepc; HEPC; LEAP1; LEAP-1; Hepcidin; Liver-expressed antimicrobial peptide 1; CNCM I-3794

**UniProt Accession Number of Target Protein:** P81172

**Published Application(s):** functional assay, WB, ELISA, IHC

**Published Species Reactivity:** Human

**Immunogen:** The original antibody was generated by immunizing mice with human hepcidin peptide 'DTHFPICIFCCGCCHRSKCGMCKT'.

**Specificity:** This antibody is specific for amino acids 'DTHFPICIFCCGCCHRSKCGMCKT' of human hepcidin-25 and does not cross react with murine hepcidin-25. Hepcidin is a an iron-regulating peptide hormone made in the liver. Hepcidin is a cationic, cysteine-rich and tightly folded peptide stabilized by 4 disulfide bonds that plays a major role in innate immunity and iron homeostasis. It controls the delivery of iron to blood plasma from intestinal cells absorbing iron, from erythrocyte-recycling macrophages, and from iron-storing hepatocytes. Hepcidin acts by binding to and inactivating the sole cellular iron exporter, ferroprotein, which delivers iron to plasma from all iron-transporting cells.

**Application Notes:** The original mouse IgG1 antibody binds human hepcidin-25 with a binding affinity of  $K_d = 99$  pM. The binding characterization of this antibody to synthetic human hepcidin-25 was done using ELISA. This antibody can be used for the identification of synthetic human hepcidin or human hepcidin from the sera of patients with an inflammatory disease using dot blot and western blot. This antibody was also used for the immunohistochemical analysis of paraffine-embedded human liver biopsies. The biological activity of this antibody was tested by incubating this antibody with macrophages from J774 murine cell line, and it was reported that this antibody induces ferroprotein degradation (US8487081).

**Antibody First Published in:** [PMID:](#)

**Note on publication:**

### Product Form

**Size:** 1 mg Purified antibody in bulk size.

**Purification:** Protein A affinity purified

**Supplied In:** PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without

added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.