

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



Anti-spike protein [PZZ 24] , 500  $\mu g,$  Ab03631-30.11-BT View online

## Anti-spike protein [PZZ 24] Ab03631-30.11-BT

This is an scFv fragment with a His tag.

Isotype and Format: scFv fragment (His), ScFv

Clone Number: PZZ 24

Alternative Name(s) of Target: Spike glycoprotein; S glycoprotein; E2; Peplomer protein

UniProt Accession Number of Target Protein:  $\ensuremath{\texttt{Q91AV1}}$ 

Published Application(s): in vivo protection, ELISA, IF

Published Species Reactivity: Porcine epidemic diarrhea virus

**Immunogen:** A scFv antibody phage display library was constructed from peripheral blood lymphocytes of piglets induced with PEDV. The library was screened with four rounds of biopanning using purified PEDV antigen, and scFv antibodies that bound to PEDV were obtained.

**Specificity:** The antibody binds the S1 region of the spike protein of PEDV. The S protein is a type I glycoprotein that plays a crucial role in virus attachment, entry, receptor binding, cell membrane fusion and induction of neutralizing antibodies. The S protein can be cleaved into S1 (residues 1–789) and S2 subunits (residues 790–1386) by host protease. The S1 subunit contains the N-terminal domain (NTD, residues 1–233) that shows sialic acid binding activity and the C-terminal domain (CTD, residues 253–638) that attaches to the cell surface receptor (e.g., aminopeptidase N (APN)). The S2 subunit mediates virus-cell membrane fusion.

**Application Notes:** The specificity of the original format of the antibody was confirmed by ELISA analysis. Immunofluorescence assay revealed that the scFv stained PEDV-infected Vero E6 cells but did not react with uninfected control cells. The scFv was shown to inhibit PEDV infectivity by the plaque reduction neutralization assay (virus neutralization titer =  $6.25 \mu g/mL$ ). Results showed the a cocktail of the scF fragment, PZZ 21 and PZZ 35 neutralized viral infection at  $3.125 \mu g/mL$ . Finally, piglets orally administered with a mixture of scFv fragment, PZZ 21 and PZZ 35 showed no to mild clinical symptoms, significantly less viral shedding, no mortality and no intestinal lesions.

**Antibody First Published in:** Zhang et al. Single Chain Fragment Variable (scFv) Antibodies Targeting the Spike Protein of Porcine Epidemic Diarrhea Virus Provide Protection against Viral Infection in Piglets. Viruses. 2019 Jan; 11(1): 58. PMID:30646521

**Note on publication:** The original paper describes the generation and characterization of the antibody.

### **Product Form**

Size:

500 μg Purified antibody in bulk size.
Purification: Purified by Immobilized Metal Affinity Chromatography
Supplied In: PBS only.
Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without

added preservatives, it is therefore recommed this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

#### Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.