



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-IL-33 [309h] Standard Size Ab03652-3.0

Isotype and Format: Mouse IgG2b, Kappa

Clone Number: 309h

Alternative Name(s) of Target: Interleukin-33; C9orf26; IL1F11; NFHEV; Interleukin-1 family member 11; IL-1F11; Nuclear factor from high endothelial venules; NF-HEV

UniProt Accession Number of Target Protein: O95760

Published Application(s): functional assay, inhibit, Block, ELISA

Published Species Reactivity: Human

Immunogen: The original mouse antibody was generated by immunizing BALB/c mice with recombinant human IL-33 (112-270 amino acid fragment)-his tagged protein. The humanized version was generated by grafting CRDs of mouse antibody onto human framework regions.

Specificity: This antibody binds an epitope between amino acid residues Ser112-Thr270 of the human IL-33. Interleukin 33 (IL-33) is a multifunctional cytokine which is constitutively expressed in structural cells such as smooth muscle cells, fibroblasts, mast cells, dendritic cells, macrophages, osteoblasts, epithelial cells, and endothelial cells. It is involved in the maturation of Th2 cells inducing the secretion of T-helper type 2-associated cytokines like IL-4, IL-5 and IL-13. It is also involved in activation of mast cells, basophils, eosinophils and natural killer cells. A study found that IL-33 is a bifunctional protein. On the one hand, IL-33 is localized in the nucleus and acts as a transcription factor; on the other hand, IL-33 is secreted outside the cell and acts as a cytokine by interacting with its receptor ST2.

Application Notes: The humanized IgG1 version of this antibody binds human IL-33 and blocks the binding of human IL-33 to IL-33R-ECD-HFC with an IC50 of 65.93ng/ml. The binding characterization of this antibody to human IL-33 was done using ELISA. This antibody was reported to inhibit IL-33 induced secretion of IL-5 by KU812 cells (CN112480252).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.