



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-GDF15 [MBM-14] Standard Size Ab03702-3.0

Isotype and Format: Mouse IgG2b, Kappa

Clone Number: MBM-14

Alternative Name(s) of Target: MIC-1; GDF-15 MIC1; PDF; PLAB; PTGFB; NAG-1; NRG-1; NAG1; NRG1; Growth/differentiation factor 15; growth and differentiation factor 15; Macrophage inhibitory cytokine 1; NSAID-activated gene 1 protein; NSAID-regulated gene 1 protein; Placental TGF-beta; Placental bone morphogenetic protein; Prostate differentiation factor

UniProt Accession Number of Target Protein: Q99988

Published Application(s): functional assay, inhibit, ELISA

Published Species Reactivity: Human

Immunogen: This antibody was isolated from a human single chain Fv (scFv) phage library using phage display technology. The library was generated from the mRNA obtained from the spleen of an immunized mouse injected with recombinant human GDF-15 protein.

Specificity: This antibody binds the human growth/differentiation factor 15 (GDF-15) also known as macrophage inhibitory cytokine 1. GDF-15 is a divergent member of the transforming growth factor- β (TGF- β) superfamily. While physiological expression is barely detectable in most somatic tissues in humans, GDF-15 is abundant in placenta. It regulates food intake, energy expenditure and body weight in response to metabolic and toxin-induced stresses. Highly elevated GDF-15 levels are mostly linked to pathological conditions including inflammation, myocardial ischemia, and notably cancer. GDF-15 is widely explored as a biomarker for disease prognosis.

Application Notes: The binding characterization of this antibody was done using ELISA. This antibody was reported to inhibit tube formation of HUVEC cells and branching of arterial sections. This antibody can effectively block the angiogenesis-inducing function of MIC-1 (KR101346132).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.