



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-IL-4 [25D2-9] Bulk Size Ab03715-10.3-BT

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This is a reformatted human IgG1 Fc Silent Fc Silent™ antibody, based on the original human IgG1 format, created for improved compatibility with existing reagents, assays and techniques.

**Isotype and Format:** Human IgG1, Fc Silent™, Kappa

**Clone Number:** 25D2-9

**Alternative Name(s) of Target:** IL4; BSF-1; Interleukin-4; B-cell stimulatory factor 1; Binetrakin; Lymphocyte stimulatory factor 1; Pitrakinra; MP4.25D2.11; ATCC HB 9809

**UniProt Accession Number of Target Protein:** P05112

**Published Application(s):** inhibit, ELISA

**Published Species Reactivity:** Human

**Immunogen:** The original antibody was generated by immunizing Lewis rat with 1ml of human IL-4. The humanized version of the antibody was created later by grafting CDRs from the original rat antibody onto human framework regions.

**Specificity:** This antibody binds human IL-4. Interleukin-4 is a cytokine secreted primarily by mast cells, T-cells, eosinophils, and basophils that plays a role in regulating antibody production, hematopoiesis and inflammation, and the development of effector T-cell responses. It induces differentiation of naive helper T cells (Th0 cells) to Th2 cells. It also induces expression of class II MHC molecules on resting B-cells and enhances both secretion and cell surface expression of IgE and IgG1.

**Application Notes:** The binding characterization of this antibody for human IL-4 was done using ELISA. This antibody was also capable of inhibiting binding of radiolabeled hIL-4 to recombinant hIL-4 receptors expressed on the Jijoye CJ cell line. This antibody was also reported to inhibit induction of human growth hormone in the germline epsilon reporter assay (US5597710; US5863537).

**Antibody First Published in:** [PMID:](#)

**Note on publication:**

## Product Form

**Size:** 1 mg Purified antibody in bulk size.

**Purification:**

Protein A affinity purified

**Supplied In:** PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.