

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

#### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



Anti-IL-4 [25D2-9] Bulk Size, 1 mg, Ab03715-3.3-BT View online

## Anti-IL-4 [25D2-9] Bulk Size Ab03715-3.3-BT

This antibody was created using our proprietary Fc Silent<sup>™</sup> engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This is a reformatted mouse IgG2b Fc Silent<sup>™</sup> antibody, based on the original mouse IgG1 format, created for improved compatibility with existing reagents, assays and techniques.

**Isotype and Format:** Mouse IgG2b, Fc Silent<sup>™</sup>, Kappa

Clone Number: 25D2-9

**Alternative Name(s) of Target:** IL4; BSF-1; Interleukin-4; B-cell stimulatory factor 1; Binetrakin; Lymphocyte stimulatory factor 1; Pitrakinra; MP4.25D2.11; ATCC HB 9809

UniProt Accession Number of Target Protein: P05112

Published Application(s): inhibit, ELISA

Published Species Reactivity: Human

**Immunogen:** The original antibody was generated by immunizing Lewis rat with 1ml of human IL-4. The humanized version of the antibody was created later by grafting CDRs from the original rat antibody onto human framework regions.

**Specificity:** This antibody binds human IL-4. Interleukin-4 is a cytokine secreted primarily by mast cells, Tcells, eosinophils, and basophils that plays a role in regulating antibody production, hematopoiesis and inflammation, and the development of effector T-cell responses. It induces differentiation of naive helper T cells (Th0 cells) to Th2 cells. It also induces expression of class II MHC molecules on resting B-cells and enhances both secretion and cell surface expression of IgE and IgG1.

**Application Notes:** The binding characterization of this antibody for human IL-4 was done using ELISA. This antibody was also capable of inhibiting binding of radiolabeled hIL-4 to recombinant hIL-4 receptors expressed on the Jijoye CJ cell line. This antibody was also reported to inhibit induction of human growth hormone in the germline epsilon reporter assay (US5597710; US5863537).

Antibody First Published in: PMID: Note on publication:

### **Product Form**

Size: 1 mg Purified antibody in bulk size. Purification: Protein A affinity purified **Supplied In:** PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommed this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.