



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-CD112 [Nec1-554-1] Standard Size Ab03744-10.3

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

**Isotype and Format:** Human IgG1, Fc Silent™, Kappa

**Clone Number:** Nec1-554-1

**Alternative Name(s) of Target:** Nectin-2; NECTIN2; HVEB; PRR2; PVRL2; Herpes virus entry mediator B; Herpesvirus entry mediator B; Nectin cell adhesion molecule 2; Poliovirus receptor-related protein 2

**UniProt Accession Number of Target Protein:** Q92692

**Published Application(s):** functional assay, inhibit, ELISA, FC

**Published Species Reactivity:** Human

**Immunogen:** The original antibody was generated by immunizing KM mice with an emulsion prepared by mixing the Nectin-2ED-Fc protein and Nectin-2ED-FLAG protein.

**Specificity:** This antibody binds an epitope between amino acids 175-240 of the human CD112 (Nectin-2). It cross reacts with cynomolgus monkey CD112. It is also capable of binding AN77-78PD, G113R and H128R mutants of human CD112. Nectin-2 acts as a modulator of T-cell signaling. It can be either a costimulator of T-cell function, or a coinhibitor, depending on the receptor it binds to. Upon binding to CD226, stimulates T-cell proliferation and cytokine production, including that of IL2, IL5, IL10, IL13, and IFNG. Upon interaction with PVRIG, inhibits T-cell proliferation.

**Application Notes:** The binding of this antibody to CD112 or Nectin-2 expressed on the surface of CHO cells was determined using whole cell ELISA. The cross reactivity of this antibody for cynomolgus monkey CD112 expressed on CHO-K1 cells was checked using flow cytometry. This antibody shows good ADCC activity against OV-90, MDA-MB-231, NCI-H1299, Du145 and U937 cell lines. The in-vivo anti-tumor activity of this antibody was tested in nude mice models transplanted with OV-90 human ovarian cancer cell line. This antibody showed T/C (Treatment/Control) value of 34%. The in-vivo activity was also checked in SCID mouse models for treating lung metastasis with MDA-MB-231 human breast cancer cell line (US20100008928).

**Antibody First Published in:** [PMID:](#)

**Note on publication:**

### Product Form

**Size:** 100 µg Purified antibody.

**Purification:**

Protein A affinity purified

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.