



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-CD112 [Nec8-4116-8] Bulk Size Ab03746-10.3-BT

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This is a reformatted human IgG1 Fc Silent Fc Silent™ antibody, based on the original human IgG1 format, created for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Human IgG1, Fc Silent™, Kappa

Clone Number: Nec8-4116-8

Alternative Name(s) of Target: Nectin-2; NECTIN2; HVEB; PRR2; PVRL2; Herpes virus entry mediator B; Herpesvirus entry mediator B; Nectin cell adhesion molecule 2; Poliovirus receptor-related protein 2

UniProt Accession Number of Target Protein: Q92692

Published Application(s): functional assay, inhibit, ELISA, FC

Published Species Reactivity: Human

Immunogen: The original antibody was generated by immunizing KM mice with an emulsion prepared by mixing the Nectin-2ED-Fc protein and Nectin-2ED-FLAG protein.

Specificity: This antibody binds human CD112 (Nectin-2) and cross reacts with cynomolgus monkey CD112. It is also capable of binding AN77-78PD, G113R and H128R mutants of human CD112. Nectin-2 acts as a modulator of T-cell signaling. It can be either a costimulator of T-cell function, or a coinhibitor, depending on the receptor it binds to. Upon binding to CD226, stimulates T-cell proliferation and cytokine production, including that of IL2, IL5, IL10, IL13, and IFNG. Upon interaction with PVRL2, inhibits T-cell proliferation.

Application Notes: The binding of this antibody to CD112 or Nectin-2 expressed on the surface of CHO cells was determined using whole cell ELISA. The cross reactivity of this antibody for cynomolgus monkey CD112 expressed on CHO-K1 cells was checked using flow cytometry. This antibody shows good ADCC activity against OV-90, MDA-MB-231, NCI-H1299, Du145 and U937 cell lines. The in-vivo activity of this antibody was checked in SCID mouse models for treating lung metastasis with MDA-MB-231 human breast cancer cell line and nude mice models for treating liver metastasis with OV-90 human ovarian cancer cell line (US20100008928).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 1 mg Purified antibody in bulk size.

Purification: Protein A affinity purified

Supplied In: PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.