



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-Cocaine [15A10] Bulk Size Ab03749-10.3-BT

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This is a reformatted human IgG1 Fc Silent Fc Silent™ antibody, based on the original human IgG1 format, created for improved compatibility with existing reagents, assays and techniques.

**Isotype and Format:** Human IgG1, Fc Silent™, Lambda

**Clone Number:** 15A10

**Alternative Name(s) of Target:** Benzoylmethylecgonine; coca; coke; crack; ChEBI: 27958

**UniProt Accession Number of Target Protein:**

**Published Application(s):** in vivo, ELISA

**Published Species Reactivity:** Species independent

**Immunogen:** The original antibody was generated by immunizing BALB/c mice with BSA conjugated transition state analogs of cocaine.

**Specificity:** This antibody hydrolyzes the benzoyl ester of cocaine to form the nonpsychoactive metabolites benzoic acid and ecgonine methylester. Cocaine is a strong stimulant used clinically as a local analgesic and vasoconstrictor and illegally as a recreational drug.

**Application Notes:** This catalytic antibody is capable of cocaine to its non psychoactive metabolites. The rate of acceleration of the catalytic antibody was reported to be ( $k_{cat}/k_{uncat} = 2.3 \times 10^4$ ). Competitive ELISA was used for the detection of binding of this antibody to the transition-state analogue of cocaine (Yang et al., 1996). In a model of cocaine overdose, this antibody protected rats from cocaine-induced seizures and sudden death in a dose-dependent fashion; a noncatalytic anticocaine antibody did not reduce toxicity. In a model of cocaine addiction, this antibody blocked completely the reinforcing effect of cocaine in rats (PMID: 9707620).

**Antibody First Published in:** Landry et al. Anti-cocaine catalytic antibodies--a novel approach to the problem of addiction. J Addict Dis. 1997;16(3):1-17. [PMID:9243335](#)

**Note on publication:** Describes the generation of catalytic antibodies with hydrolytic activity against cocaine.

## Product Form

**Size:**

1 mg Purified antibody in bulk size.

**Purification:** Protein A affinity purified

**Supplied In:** PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.