

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



## Anti-Morphine [M86] Bulk Size Ab03754-30.11-BT

This is an scFv fragment with a His tag.

This is a reformatted mouse scFv antibody, based on the original mouse scFv format, created for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: scFv fragment (His), ScFv

Clone Number: M86

Alternative Name(s) of Target: Morphium; Codeine; Heroin; Opium; ChEBI: 17303; FERM ABP-11378 UniProt Accession Number of Target Protein:

Published Application(s): ELISA

Published Species Reactivity: Species independent

**Immunogen:** The original antibody was generated by immunizing BALB/c mice with morphine-BSA conjugated fusion protein. A mouse scFv library was generated from the spleen cells and the particular antibody was isolated by panning against morphine-conjugated thyroglobulin (morphine-C-Tg). **Specificity:** This antibody binds morphine and cross reacts with other opioids like opium, codeine and heroin. Morphine is a strong opiate that is found naturally in opium, a dark brown resin in poppies (Papaver somniferum). It is considered the classic opioid analgesic, most commonly used in pain management, morphine provides major relief to patients afflicted with pain. It is a non-synthetic narcotic with a high potential for abuse.

**Application Notes:** The binding reactivity of this antibody for morphine conjugated thyroglobulin (morphine-C-Tg) was done using ELISA. The original scFv antibody binds morphine-C-Tg with a binding affinity of Kd= 0.126 nM. It was reported that this antibody does not cross react with thyroglobulin conjugated codeine-, cocaine-, or ketamine. However, this antibody is capable of cross reacting with free morphine, opium, codeine and heroin. The IC50 values of M86 scFv for opium, morphine, codeine and heroin was 257 ng/mL, 36.4, 7.3 and 7.4 nM, respectively (Matsukizono et al., 2013; JP2012232948).

**Antibody First Published in:** Matsukizono et al. Characterization of a Single Chain Fv Antibody that Reacts with Free Morphine. Antibodies 2013, 2(1), 93-112. PMID:

**Note on publication:** This paper describes the generation of and characterization of an scFv antibody against morphine.

### **Product Form**

**Size:** 500 μg Purified antibody in bulk size. **Purification:** Purified by Immobilized Metal Affinity Chromatography **Supplied In:** PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommed this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

#### Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.