



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-IDH1 R132H [10G5-1] Standard Size Ab03872-10.3

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This is a reformatted human IgG1 Fc Silent Fc Silent™ antibody, based on the original human IgG1 format, created for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Human IgG1, Fc Silent™, Kappa

Clone Number: 10G5-1

Alternative Name(s) of Target: IDH; IDH1; IDPc; Isocitrate dehydrogenase [NADP] cytoplasmic; Isocitrate dehydrogenase 1; Cytosolic NADP-isocitrate dehydrogenase; NADP(+)-specific ICDH; Oxalosuccinate decarboxylase

UniProt Accession Number of Target Protein: O75874

Published Application(s): ELISA, IHC

Published Species Reactivity: Human

Immunogen: The original antibody was generated by immunizing BALB/c mice with human IDH-R132H polypeptide (KPIIGHHAYGDQYC) with an N-terminal BSA conjugation.

Specificity: This antibody binds human isocitrate dehydrogenase 1 (IDH1) having R132H mutation. IDH1 is an enzyme that catalyzes the NADP⁺-dependent oxidative decarboxylation of isocitrate (D-threo-isocitrate) to 2-ketoglutarate (2-oxoglutarate), which is required by other enzymes such as the phytanoyl-CoA dioxygenase. It plays a critical role in the generation of NADPH, an important cofactor in many biosynthesis pathways. The NADPH produced from isocitrate dehydrogenase 1 is involved in the breakdown of fats for energy, and it also protects cells from potentially harmful molecules called reactive oxygen species. Mutation in IDH1 are found in several genetic conditions and in some types of cancer, including gliomas, cartilaginous tumors, acute myelogenous leukemia, myelodysplastic syndromes, and brain cancer.

Application Notes: The binding reactivity of this antibody to R132H mutant of human IDH1 was determined using ELISA. The original mouse IgG antibody binds IDH1 R132H mutant with an affinity of $K_d = 5.76 \times 10^{-10}$ M. This antibody was used in the immunohistochemical analysis for detection of IDH1 R132H protein in cancerous tissues (CN108727500).

Antibody First Published in: PMID:

Note on publication:

Product Form

Size: 100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.