



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-CD47 [105D11] Standard Size Ab03948-23.0

**Isotype and Format:** Rabbit IgG, Kappa

**Clone Number:** 105D11

**Alternative Name(s) of Target:** IAP; MER6; OV-3; Leukocyte surface antigen CD47; Antigenic surface determinant protein OA3; Integrin-associated protein; Protein MER6; ovarian cancer marker

**UniProt Accession Number of Target Protein:** Q08722

**Published Application(s):** Block, ELISA, FC

**Published Species Reactivity:** Human, Cynomolgus Monkey

**Immunogen:** The original antibody was generated by immunizing BALB/c mice with recombinant human CD47-mFc fusion protein.

**Specificity:** This antibody binds the extracellular domain of the human CD47 and cross reacts with cynomolgus monkey CD47. CD47 is broadly distributed on normal adult tissues, as well as ovarian tumors, and is especially abundant in some epithelia and the brain. IAP or integrin associated protein has a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins. It further plays an important role in memory formation and synaptic plasticity in the hippocampus. It also acts as a receptor for SIRPA. Binding to SIRPA prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells.

**Application Notes:** The binding characterization of this antibody to recombinant human Cd47-mFc fusion protein was done using ELISA. This antibody could also bind human and cynomolgus monkey CD47 expressed on the surface of cells in a flow cytometric assay. This antibody was also reported to block the binding between SIRPα-hFc protein and human CD47 expressed on the cell surface (CN110724672).

**Antibody First Published in:** [PMID:](#)

**Note on publication:**

### Product Form

**Size:** 100 µg Purified antibody.

**Purification:** Protein A affinity purified

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic

procedures for humans or animals.