



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-von Willebrand factor [82D6A3] Standard Size Ab03983-1.1

Isotype and Format: Mouse IgG1, Kappa

Clone Number: 82D6A3

Alternative Name(s) of Target: human vWF A3 domain; VWF-A3; vWF

UniProt Accession Number of Target Protein: P04275

Published Application(s): crystallization, in vivo, inhibition

Published Species Reactivity: Baboon, Human

Immunogen: The original antibody was purified from murine ascites by Protein A chromatography.

Specificity: This antibody is specific for human vWF, A3 domain.

Application Notes: This antibody was used as an inhibitor to study platelet adhesion to collagen VI and ECM. It was found to inhibit the binding of von Willebrand factor (VWF) to collagen I and III (Hoylaerts et al., 1997; PMID: 9164855). This antibody was found to have a comparable affinity for baboon vWF as for human vWF and inhibited the binding of baboon vWF to collagen with an inhibitory concentration of 50% (IC₅₀) of 3.5 µg/ml. Furthermore, it prevented the binding of VWF to collagen and removed bound VWF from a collagen surface, albeit with a lower efficacy. Moreover, the antithrombotic efficacy of the antibody was evaluated in a modified Folts model in baboons; this antibody was used to inhibit the interaction between collagen and vWF to prevent arterial thrombosis in baboons (Wu et al., 2002; PMID: 11986216). The crystal structure of the antibody's Fab fragment, in complex with the vWF A3 domain, was determined (Staelens et al., 2006; PMID: 16314412).

Antibody First Published in: Hoylaerts et al. von Willebrand factor binds to native collagen VI primarily via its A1 domain *Biochem J.* 1997 May 15;324 (Pt 1)(Pt 1):185-91. doi: 10.1042/bj3240185 PMID:9164855

Note on publication: The original publication investigates the interaction mechanism between collagen VI and von Willebrand factor (vWF).

Product Form

Size: 100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration:

1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.