



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Gamma Delta TCR [G5-4] Bulk Size Ab03990-2.0-BT

Isotype and Format: Mouse IgG2a, Kappa

Clone Number: G5-4

Alternative Name(s) of Target: TCR $\gamma\delta$; $\gamma\delta$ TCR; TCR GD; GD TCR; TCRg/d; T-cell surface glycoprotein CD3 delta chain; CD3D; T3D; T-cell receptor T3 delta chain; T-cell surface glycoprotein CD3 gamma chain; CD3G; T-cell receptor T3 gamma chain; CD3g; T3G

UniProt Accession Number of Target Protein: P04234; P09693

Published Application(s): functional assay, FC

Published Species Reactivity: Human

Immunogen: The original antibody was generated by immunizing BALB/c mice with recombinant $\gamma\delta$ (OT3)-Fc fusion protein as an immunogen.

Specificity: This antibody is specific for the human TCR delta and gamma chains. The T-cell receptor (TCR) is a molecule found on the surface of T cells, or T lymphocytes, that is responsible for recognizing fragments of antigen as peptides bound to major histocompatibility complex (MHC) molecules. On activation through antigen presenting cells (APCs), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. TCR part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response.

Application Notes: The binding of this antibody to $\gamma\delta$ T cells was determined using flow cytometry. This antibody can compete with commercially available anti- $\gamma\delta$ TCR antibody IMMU510 for binding to gamma-delta T cells. It is proposed that, since there is some competition between the two antibodies, G5-4 binds a similar or identical epitope to IMMU510. In an in vitro proliferation assay, this antibody was reported to positively stimulate the proliferation of $\gamma\delta$ T cells. The scFv version of this antibody was also reported to stimulate the secretion of cytokines like IFN- γ and TGF- α . G5-4ScFv-amplified $\gamma\delta$ T cells can effectively kill daudi cells, and the killing efficiency is greater than 85% when the effect-to-target ratio is 5:1 (CN103130894).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 1 mg Purified antibody in bulk size.

Purification: Protein A affinity purified

Supplied In:

PBS only.

Storage Recommendation: Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.